

Scalar Potential Without Cubic Term in 3-3-1 Models Without Exotic Electric Charges

Yithsbey Giraldo^{1,2} and William A. Ponce²

¹ Departamento de Física, Universidad de Nariño, A.A. 1175, Pasto, Colombia.

² Instituto de Física, Universidad de Antioquia, A.A. 1226, Medellín, Colombia.

Received: date / Revised version: date

Abstract. A detailed study of the criteria for stability of the scalar potential, and the proper electroweak symmetry breaking pattern in some 3-3-1 models without exotic electric charges is presented. In this paper we concentrate in a scalar sector with three Higgs scalar triplets, with a potential that does not include the cubic term, due to the presence of a discrete symmetry. For the analysis we use, and improve, a method previously developed to study the scalar potential in the two-Higgs-doublet extension of the standard model. Our main result is to show the consistency of those 3-3-1 models without exotic electric charges.

1 Introduction

A simple extension of the standard model (SM) consists of adding to the model a second Higgs scalar doublet [1], defining in this way the so-called two Higgs doublet model (THDM). The different ways how the two Higgs scalar doublets couple to the fermion sector, define the different versions of this extension[1,2]. Many gauge group extensions of the SM have the THDM as an effective low energy theory (in this regard see the papers in [2] and references therein). In these extensions one intermediate step in the symmetry breaking chain leads to the $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ gauge theory with two Higgs doublets in one of its several versions.

A novel method for a detailed analysis of the scalar potential in the most general THDM was presented in Refs. [3,4] where by using powerful algebraic techniques, the authors studied in detail the stationary points of the scalar potential. This allowed them to give, in a very concise way, clear criteria for the stability of the scalar potential and for the correct electroweak symmetry breaking pattern. By using different approaches, the authors in Refs. [5] reached also interesting conclusions for the scalar potential of the THDM, some of them related to the ones presented in Ref.[3].

In the present work we make use of some of the new algebraic developments cited in the former paragraph, to analyze the scalar sector of an extension to the SM based on the local gauge group $SU(3)_c \otimes SU(3)_L \otimes U(1)_X$ [6,7,8,9,10] (called hereafter 3-3-1 for short).

In general, the scalar sector for 3-3-1 models is quite complicated and difficult to analyze in detail. For example, for the minimal model (the Pisano-Pleitez-Frampton model [6]) three Higgs scalar triplets and one additional Higgs sextet must be used, in order to break the symme-

try and provide, at the same time, masses to the fermion fields. For the 3-3-1 models without exotic electric charges [7,8,9,10] the situation is simpler because it turns out that less Higgs scalar multiplets are needed [11]. For example, the so called economical 3-3-1 model [12] makes use of only two Higgs scalar triplets which are able to break the symmetry in a consistent way, although they are not able to produce a consistent fermion mass spectrum at tree-level. The alternative approach is to deal with three Higgs scalar triplets instead of two, as done for example in Refs. [7,8].

In this work we pursue the study of the scalar sector of the 3-3-1 models without exotic electric charges started in Ref. [13], by considering this time a model with three Higgs scalar triplets. A discrete symmetry will be applied to the corresponding scalar potential [14,15], which simplifies and facilitates its analysis, due that the cubic (or trilinear) term would be absent. In this analysis we will derive constraints on the parameters of the scalar potential coming from its stability and from the electroweak symmetry breaking conditions (the stability of an scalar potential at the classical level, which is fulfilled when it is bounded from below, is a necessary condition in order to have a consistent theory). The global minimum of the potential will also be found by determining its stationary points.

This paper is organized as follows: in Sect. 2 we introduce the 3-3-1 models without exotic electric charges and review the different scalar sectors available in the literature for this type of models; in Sect. 3 we introduce the scalar potential under study and analyze the consistency of the electroweak symmetry breaking pattern proposed; then in Sect. 4 we study the stability of the scalar potential, and in Sect. 5 we find its stationary points and its global minimum. Our conclusions are presented in Sect. 6.

Three appendixes, one with a general review and two technical ones are presented at the end.

2 3-3-1 models without exotic electric charges

As demonstrated in Refs. [9, 10, 12], there exist a total of eight different three-family 3-3-1 models without exotic electric charges, that do not contain fermion singlets or in vector-like representations. Each model has a different spin 1/2 particle structure, but they have the same gauge boson sector. In principle, all can bear the same scalar sector too.

2.1 The Gauge sector

The gauge boson structure for any 3-3-1 model without exotic electric charges is: one gauge field B^μ associated with $U(1)_X$, the 8 gluon fields G^μ associated with $SU(3)_c$ which remain massless after breaking the symmetry, and 8 gauge fields from $SU(3)_L$ that we write as [12]

$$\frac{1}{\sqrt{2}}\lambda_\alpha A_\alpha^\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} D_1^\mu & W^{+\mu} & K^{+\mu} \\ W^{-\mu} & D_2^\mu & K^{0\mu} \\ K^{-\mu} & \bar{K}^{0\mu} & D_3^\mu \end{pmatrix},$$

where $D_1^\mu = A_3^\mu/\sqrt{2} + A_8^\mu/\sqrt{6}$, $D_2^\mu = -A_3^\mu/\sqrt{2} + A_8^\mu/\sqrt{6}$, and $D_3^\mu = -2A_8^\mu/\sqrt{6}$. λ_i , $i = 1, 2, \dots, 8$, are the eight Gell-Mann matrices normalized as $Tr(\lambda_i \lambda_j) = 2\delta_{ij}$.

The charge operator associated with the unbroken gauge symmetry $U(1)_Q$ is

$$Q = \frac{\lambda_3}{2} + \frac{\lambda_8}{2\sqrt{3}} + XI_3 \quad (1)$$

where $I_3 = \text{Diag.}(1, 1, 1)$ is the diagonal 3×3 unit matrix, and the X values are related to the $U(1)_X$ hypercharge and are fixed by anomaly cancellation. The sine square of the electroweak mixing angle is given by $S_W^2 = 3g_1^2/(3g_3^2 + 4g_1^2)$ where g_1 and g_3 are the coupling constants of $U(1)_X$ and $SU(3)_L$ respectively. The photon field is given by

$$A_0^\mu = S_W A_3^\mu + C_W \left[\frac{T_W}{\sqrt{3}} A_8^\mu + \sqrt{(1 - T_W^2/3)} B^\mu \right], \quad (2)$$

where C_W and T_W are the cosine and tangent of the electroweak mixing angle, respectively.

There are two weak neutral currents in the model associated with the two flavor diagonal neutral gauge weak bosons

$$Z_0^\mu = C_W A_3^\mu - S_W \left[\frac{T_W}{\sqrt{3}} A_8^\mu + \sqrt{(1 - T_W^2/3)} B^\mu \right],$$

$$Z_0^{\prime\mu} = -\sqrt{(1 - T_W^2/3)} A_8^\mu + \frac{T_W}{\sqrt{3}} B^\mu, \quad (3)$$

and one current associated with the flavor non diagonal neutral gauge boson $K^{0\mu}$ which is charged in the sense that it has a kind of weak V isospin charge. In the former expressions Z_0^μ coincides with the weak neutral current of the SM.

2.2 A Fermion sector

The particular 3-3-1 model without exotic electric charges most extensively studied in the literature, known as the 3-3-1 model with right-handed neutrinos, has the following anomaly free fermion sector [7]:

$$\begin{aligned} \psi_L^a &= (l^{-a}, \nu^a, N^{0a})_L^T \sim (1, 3^*, -1/3), \\ l_L^{+a} &\sim (1, 1, 1), \\ Q_L^i &= (u^i, d^i, D^i)_L^T \sim (3, 3, 0), \\ Q_L^3 &= (d^3, u^3, U)_L^T \sim (3, 3^*, 1/3), \\ u_L^{ca} &\sim (3^*, 1, -2/3), \quad d_L^{ca} \sim (3^*, 1, 1/3), \\ U_L^c &\sim (3^*, 1, -2/3), \quad D_L^{ci} \sim (3^*, 1, 1/3), \end{aligned}$$

where the numbers inside the parenthesis stand for $[SU(3)_c, SU(3)_L, U(1)_Y]$ representations, $a = 1, 2, 3$ is a family index and $i = 1, 2$ is related to two of the three families. D^i and U are three exotic quarks with electric charges $-1/3$, $-1/3$ and $2/3$, respectively, and N_L^{0a} can play the role of right-handed neutrinos. The three lepton families are arranged in antitriplets of $SU(3)_L$. In order to cancel the $SU(3)_L$ anomaly, two quark families must transform as $SU(3)_L$ triplets and the remaining one as an antitriplet. It is customary to arrange the first two quark families in triplets and the third one in an antitriplet. This choice is meant to distinguish the possible new dynamics effects arising in the third family.

2.3 The scalar sector

As far as we know, for the 3-3-1 models without exotic electric charges, three different scalar sectors have been used in the literature, to deal with the spontaneous breaking of the gauge symmetry down to $U(1)_Q$ and, to produce at the same time, masses for the Fermion fields. Each set, as described anon, has its own advantages and disadvantages. They are:

2.3.1 The economical model

Introduced in the literature in Ref. [12] and further analyzed in Refs. [16, 17]. It makes use of only two scalar triplets, which together with their vacuum expectation values (VEV) are:

$$\Phi_1(1, 3^*, -1/3) = \begin{pmatrix} \phi_1^- \\ \phi_1^0 \\ \phi_1^{0'} \end{pmatrix}, \quad \text{with VEV: } \langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \\ V_1 \end{pmatrix}, \quad (4a)$$

$$\Phi_3(1, 3^*, 2/3) = \begin{pmatrix} \phi_3^0 \\ \phi_3^+ \\ \phi_3^{+'} \end{pmatrix}, \quad \text{with VEV: } \langle \Phi_3 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_3 \\ 0 \\ 0 \end{pmatrix}. \quad (4b)$$

The former structure is the simplest one, able to break the 3-3-1 symmetry down to $U(1)_Q$ in a consistent way [12].

In spite of its simplicity, it has the disadvantage of being unable to produce a consistent Fermion mass spectrum at tree-level. The claim in Ref. [17] is that the quantum fluctuations can generate non-zero mass terms for all the Fermion fields, but a systematic (tedious) numerical analysis reproducing the fermion mass spectrum has not been published yet, although probably, the most serious hurdle for the survival of this model, is the existence of flavor changing neutral currents (FCNC) at tree-level, mediated by the Higgs scalar fields (only two sets of scalar fields producing masses for three Fermion families), neutral currents that severely constraint the parameters of the model.

For this economical model, the study of the scalar potential, using the method introduced in Refs. [3,4] (and briefly review in appendix A) has been presented in full detail in Ref. [13].

2.3.2 The set with three scalar triplets

This set makes use of the two scalar Higgs fields of the economical model, plus the extra one

$$\Phi_2(1, 3^*, -1/3) = \begin{pmatrix} \phi_2^- \\ \phi_2^0 \\ \phi_2^0 \end{pmatrix}, \text{ with VEV: } \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 \\ V_2 \end{pmatrix}, \quad (5)$$

where we have assumed that all the five electrically neutral components do acquire non-zero VEV. This set of three scalar Higgs fields, with the vacuum aligned such that $V_1 = v_2 = 0$, was used for the first time in Refs. [7,8]. The set has the advantage of being able to produce tree-level masses for all the Fermion fields (which is true even for the particular alignment used in the original papers), but it can not completely avoid the presence of FCNC at tree-level, coming from the scalar sector.

Notice that the VEV $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ generate masses for the exotic quarks and the new heavy gauge bosons, while VEV $\langle \Phi_3 \rangle$ generates masses for ordinary fermions and for the SM gauge bosons. To keep the model consistent with low energy phenomenology, in this paper we will use $\langle \Phi_3 \rangle \neq 0$ and the hierarchy

$$V_1, V_2 \gg v_1, v_2, v_3, \quad (6)$$

except for those cases when V_1 or V_2 are zero, when the hierarchy becomes

$$V_i \gg v_1, v_2, v_3; \quad i = 1, 2. \quad (7)$$

(Taking $\langle \Phi_3 \rangle = 0$ implies that several fermion fields remain massless.)

2.3.3 The extended scalar set

Introduced in the literature in Refs. [18], it consists of four scalar triplets: Φ_1 , Φ_2 and Φ_3 as above with the vacuum

aligned such that $V_1 = v_2 = 0$ as in the original papers, plus a new scalar Higgs field

$$\Phi_4(1, 3^*, -1/3) = \begin{pmatrix} \phi_4^- \\ \phi_4^0 \\ \phi_4^0 \end{pmatrix}, \text{ with VEV: } \langle \Phi_4 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 0 \\ v_4 \end{pmatrix}, \quad (8)$$

with the hierarchy $v_1 \sim v_3 \sim v_4 \ll V_2 \sim 1$ TeV. This set of four scalar fields combined with a convenient discrete symmetry [18], is able to generate several see-saw mechanisms, the basis of a consistent Fermion mass spectrum, and avoid, at the same time, tree-level FCNC coming from the scalar sector.

In the following analysis we will concentrate only in the set with 3 scalar triplets as defined in Sects. (2.3.1) and (2.3.2), with the most general VEV structure, but with the constraint derived in Appendix B.

3 The scalar Potential

The most general scalar potential which is 3-3-1 invariant, for the set of three scalar triplets Φ_1 , Φ_2 and Φ_3 is given by

$$\begin{aligned} V'(\Phi_1, \Phi_2, \Phi_3) = & \mu_1^2 \Phi_1^\dagger \Phi_1 + \mu_2^2 \Phi_2^\dagger \Phi_2 + \mu_3^2 \Phi_3^\dagger \Phi_3 \\ & + \frac{1}{2} (\mu_4^2 \Phi_1^\dagger \Phi_2 + \mu_4^{2*} \Phi_2^\dagger \Phi_1) + \lambda_1 (\Phi_1^\dagger \Phi_1)^2 + \lambda_2 (\Phi_2^\dagger \Phi_2)^2 \\ & + \lambda_3 (\Phi_3^\dagger \Phi_3)^2 + \lambda_4 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_5 (\Phi_1^\dagger \Phi_1) (\Phi_3^\dagger \Phi_3) \\ & + \lambda_6 (\Phi_2^\dagger \Phi_2) (\Phi_3^\dagger \Phi_3) + \lambda_7 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ & + \lambda_8 (\Phi_1^\dagger \Phi_3) (\Phi_3^\dagger \Phi_1) + \lambda_9 (\Phi_2^\dagger \Phi_3) (\Phi_3^\dagger \Phi_2) \\ & + (f \Phi_1^\dagger \Phi_2 + f^* \Phi_2^\dagger \Phi_1)^2 + \frac{1}{2} (\lambda_{11} \Phi_1^\dagger \Phi_2 + \lambda_{11}^* \Phi_2^\dagger \Phi_1) (\Phi_1^\dagger \Phi_1) \\ & + \frac{1}{2} (\lambda_{12} \Phi_1^\dagger \Phi_2 + \lambda_{12}^* \Phi_2^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\ & + \frac{1}{2} (\lambda_{13} \Phi_1^\dagger \Phi_2 + \lambda_{13}^* \Phi_2^\dagger \Phi_1) (\Phi_3^\dagger \Phi_3) \\ & + \frac{1}{2} [\lambda_{14} (\Phi_1^\dagger \Phi_3) (\Phi_3^\dagger \Phi_2) + \lambda_{14}^* (\Phi_3^\dagger \Phi_1) (\Phi_2^\dagger \Phi_3)] \\ & + (g \epsilon_{ijk} \Phi_1^i \Phi_2^j \Phi_3^k + h.c.). \end{aligned} \quad (9)$$

Since μ_4^2 , f , λ_{11} , λ_{12} , λ_{13} , λ_{14} and the trilinear coupling constant g can be complex numbers, there are 26 free parameters in $V'(\Phi_1, \Phi_2, \Phi_3)$ and 5 VEV, in principle all of them different from zero. The last element of V' correspond to the so called cubic term of the potential, which is closely related to a determinant function of Higgs fields due to the Levi-Civita component ϵ_{ijk} .

For the sake of simplicity we are going to assume real VEV throughout this paper, which means that spontaneous CP violation is not going to be considered in our analysis. Notice also that the most general scalar potential $V'(\Phi_1, \Phi_2, \Phi_3)$ in (9) is invariant under the local Gauge group $SU(3)_L \otimes U(1)_X$, invariance that is spontaneously broken by the VEV in $\langle \Phi_i \rangle$, $i = 1, 2, 3$ down to $U(1)_Q$, where Q is the electric charge generator in equation (1). So, after the breaking of the symmetry, a consistent model will emerge only if eight massless Goldstone Bosons show

up, coming from the transformed potential obtained from (9); zero mass Bosons that should be eaten up by the Gauge Bosons associated with the $SU(3)_L \otimes U(1)_X$ broken symmetry.

The scalar potential in Eq. (9) is quite complicated and very difficult to study in a systematic way, and as far as we know, it has not been studied in full detail in the literature yet (and we do not intend to do it here either). A partial analysis of this general potential, for the particular vacuum alignment $V_1 = v_2 = 0$, has been done in Ref. [19]. However, as mentioned in Refs. [14,15], by introducing discrete symmetries, the form of the potential simplifies largely and can be analyzed in detail, as we are going to do next.

3.1 Discrete symmetry in the scalar potential

Under assumption of the discrete symmetry $\Phi_1 \rightarrow -\Phi_1$, the most general potential obtained from (9), is presented in Appendix C, where it is demonstrated that f can be taken as a single parameter. As a consequence of this, the reduced potential

$$\begin{aligned} V(\Phi_1, \Phi_2, \Phi_3) = & \mu_1^2 \Phi_1^\dagger \Phi_1 + \mu_2^2 \Phi_2^\dagger \Phi_2 + \mu_3^2 \Phi_3^\dagger \Phi_3 \\ & + \lambda_1 (\Phi_1^\dagger \Phi_1)^2 + \lambda_2 (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_3^\dagger \Phi_3)^2 \\ & + \lambda_4 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) + \lambda_5 (\Phi_1^\dagger \Phi_1) (\Phi_3^\dagger \Phi_3) \\ & + \lambda_6 (\Phi_2^\dagger \Phi_2) (\Phi_3^\dagger \Phi_3) + \lambda_7 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ & + \lambda_8 (\Phi_1^\dagger \Phi_3) (\Phi_3^\dagger \Phi_1) + \lambda_9 (\Phi_2^\dagger \Phi_3) (\Phi_3^\dagger \Phi_2) \\ & + \frac{\lambda_{10}}{2} (\Phi_1^\dagger \Phi_2 + \Phi_2^\dagger \Phi_1)^2, \end{aligned} \quad (10)$$

contains only 13 free parameters (instead of 26) and does not include the cubic term $\epsilon_{ijk} \Phi_1^i \Phi_2^j \Phi_3^k$. The rest of the paper will be dedicated to study this potential (10).

A careful analysis shows now that, due to the absence of the cubic term, the potential $V(\Phi_1, \Phi_2, \Phi_3)$ turns out to be $U(3) \otimes U(1)_X$ invariant [instead of $SU(3) \otimes U(1)_X$] with the consequence that the most general VEV breaks this symmetry down to $U(1)_Q$ as before, producing now nine Goldstone Bosons, instead of the eight that can be Gauged away, due to the fact that the generator $I_3 = Dg(1, 1, 1)$ gets also broken. This leaves an (unphysical?) extra zero mass scalar after the implementation of the Higgs mechanism. The simplest way to avoid this situation is by restoring the cubic term in the scalar potential (a dynamical breaking of the $U(3)$ symmetry), something does not allowed by the discrete symmetry imposed. But as shown in appendix (B), for the case when $\langle \Phi_3 \rangle \neq 0$ the problem can be solved by demanding that $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ became linearly dependent (LD); this avoids an spontaneous breaking of the $U(3) \otimes U(1)_X$ symmetry down to $SU(3) \otimes U(1)_X$, with the consequence that the VEV must satisfy the constraint

$$v_2 V_1 = v_1 V_2, \quad (11)$$

which can be used to express at least one VEV in terms of the rest.

Notice that if $\langle \Phi_3 \rangle = 0$, the $U(3)$ generator $Diag.(1, 0, 0)$ remains unbroken by $\langle \Phi_1 \rangle \oplus \langle \Phi_2 \rangle \oplus \langle \Phi_3 \rangle$, restoring in this

way the eight Goldstone Bosons required. But we are not going to consider this unphysical situation as previously mentioned.

Before continuing, let us emphasize that constraint (11) is a consequence of demanding a consistent implementation of the Higgs mechanism for the breaking of the original $SU(3) \otimes U(1)_X$ local Gauge symmetry, respecting the electromagnetic $U(1)_Q$ invariance, and it is not coming from the minimization of the scalar potential. On the contrary, this constraint is taking into account when we study the stability and minimization of the potential.

Notice that first two papers in Ref. [7] and all papers in Ref. [14], the reduced potential $V(\Phi_1, \Phi_2, \Phi_3)$ was studied using the particular vacuum alignment $V_1 = v_2 = 0$, with $V_2 \gg v_1 \neq 0$, in clear contradiction with equation (11). As an immediate consequence, in those papers an extra zero mass Goldstone Boson which cannot be Gauged away appears, making the analysis and some of the conclusions in all those papers dubious. To add in proof, notice that the four papers in Ref. [15] make use of that extra Goldstone Boson to implement the Peccei-Quinn symmetry [20] in the context of the 3-3-1 model with right handed neutrinos, with the inconvenience of having in their analysis an unrealistic axion that is hidden by the introduction of an extra scalar field.

In what follows we are going to study the consistency of the scalar potential $V(\Phi_1, \Phi_2, \Phi_3)$ in Eq. (10), under the linear dependent (LD) constraint equation: $v_2 V_1 = v_1 V_2$.

To start with, let us define as usual the scalar fields in the way:

$$\phi_1^0 = \frac{v_1 + H_1 + iA_1}{\sqrt{2}}, \quad \phi_1^{\prime 0} = \frac{V_1 + H_1' + iA_1'}{\sqrt{2}}, \quad (12a)$$

$$\phi_2^0 = \frac{v_2 + H_2 + iA_2}{\sqrt{2}}, \quad \phi_2^{\prime 0} = \frac{V_2 + H_2' + iA_2'}{\sqrt{2}}, \quad (12b)$$

$$\phi_3^0 = \frac{v_3 + H_3 + iA_3}{\sqrt{2}}, \quad (12c)$$

where a real part H is called in the literature a CP-even scalar and an imaginary part A a CP-odd scalar or pseudoscalar field.

3.2 Independent vacuum structures

Assuming for the VEV the hierarchy in (6) or in (7), and using the LD constraint relation (11), we classify in Table 1, all the possible 3-3-1 vacuum structures of Φ_1 and Φ_2 , the two scalar triplets with identical quantum numbers, where at least one VEV is different from zero.

A careful analysis shows that not all the nine structures are independent. As a matter of fact, by performing an $SU(3)_L$ transformation on $\langle \Phi_1 \rangle$ and on $\langle \Phi_2 \rangle$ in structure 1 of Table 1, we can obtain either the structure configuration 2 or the structure configuration 5. But it is not possible to make an $SU(3)_L \otimes U(1)_X$ transformation followed by a change of basis of the Higgs fields $\Phi_i \rightarrow \Phi_i'$ of the form

$$\begin{pmatrix} \Phi_1' \\ \Phi_2' \end{pmatrix} = U \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}, \quad (13)$$

Table 1. Different VEV structures for scalars Φ_1 and Φ_2 .

Structure	VEV	Vacuum alignments
1	$v_1, V_1, v_2, V_2 \neq 0$ and $v_2 V_1 = v_1 V_2$	$\langle \Phi_1 \rangle \propto (0 \ v_1 \ V_1)^T$, $\langle \Phi_2 \rangle \propto (0 \ v_2 \ V_2)^T$
2	$v_1, v_2 = 0; V_1, V_2 \neq 0$	$\langle \Phi_1 \rangle \propto (0 \ 0 \ V_1)^T$, $\langle \Phi_2 \rangle \propto (0 \ 0 \ V_2)^T$
3	$v_2, V_2 = 0; v_1, V_1 \neq 0$	$\langle \Phi_1 \rangle \propto (0 \ v_1 \ V_1)^T$, $\langle \Phi_2 \rangle = (0 \ 0 \ 0)^T$
4	$v_1, V_1 = 0; v_2, V_2 \neq 0$	$\langle \Phi_1 \rangle = (0 \ 0 \ 0)^T$, $\langle \Phi_2 \rangle \propto (0 \ v_2 \ V_2)^T$
5	$V_1, V_2 = 0; v_1, v_2 \neq 0$	$\langle \Phi_1 \rangle \propto (0 \ v_1 \ 0)^T$, $\langle \Phi_2 \rangle \propto (0 \ v_2 \ 0)^T$
6	$v_2, V_2, V_1 = 0; v_1 \neq 0$	$\langle \Phi_1 \rangle \propto (0 \ v_1 \ 0)^T$, $\langle \Phi_2 \rangle = (0 \ 0 \ 0)^T$
7	$v_2, V_2, v_1 = 0; V_1 \neq 0$	$\langle \Phi_1 \rangle \propto (0 \ 0 \ V_1)^T$, $\langle \Phi_2 \rangle = (0 \ 0 \ 0)^T$
8	$v_1, V_1, V_2 = 0; v_2 \neq 0$	$\langle \Phi_1 \rangle = (0 \ 0 \ 0)^T$, $\langle \Phi_2 \rangle \propto (0 \ v_2 \ 0)^T$
9	$v_1, V_1, v_2 = 0; V_2 \neq 0$	$\langle \Phi_1 \rangle = (0 \ 0 \ 0)^T$, $\langle \Phi_2 \rangle \propto (0 \ 0 \ V_2)^T$

where U is a 2×2 unitary matrix, such that the configuration 3 can be obtained; this is because the transformation (13) violates the discrete symmetry $\Phi_1 \rightarrow -\Phi_1$ previously imposed on the scalar potential. It is also possible to show that structures 3 and 4 are equivalent to each other due to the symmetry of the potential under the exchange $\Phi_1 \leftrightarrow \Phi_2$, with some parameters renamed appropriately. In conclusion, the analysis shows that only structures 1 and 3 in Table 1 are independent and are the only cases we are going to consider in our analysis.

3.2.1 Vacuum structure with $v_1, V_1, v_2, V_2 \neq 0$

It will be shown further below that, by minimization methods applied on potential, the scalars acquiring non-zero VEVs along their electrically neutral entries, is highly suggested (121).

$$\langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \\ V_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 \\ V_2 \end{pmatrix}, \quad \langle \Phi_3 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_3 \\ 0 \\ 0 \end{pmatrix}, \quad (14)$$

with $v_2 V_1 = v_1 V_2$. And requiring that in the shifted potential obtained from $V(\Phi_1, \Phi_2, \Phi_3)$, the linear terms in the fields must be absent, we get in the tree-level approximation the following constraint equations

$$2\mu_1^2 + (\lambda_7 + \lambda_4 + 2\lambda_{10}) (V_2^2 + v_2^2) + 2\lambda_1 (V_1^2 + v_1^2) + \lambda_5 v_3^2 = 0, \quad (15a)$$

$$2\mu_2^2 + 2\lambda_2 (V_2^2 + v_2^2) + (\lambda_7 + \lambda_4 + 2\lambda_{10}) (V_1^2 + v_1^2) + \lambda_6 v_3^2 = 0, \quad (15b)$$

$$2\mu_3^2 + \lambda_6 (V_2^2 + v_2^2) + \lambda_5 (V_1^2 + v_1^2) + 2\lambda_3 v_3^2 = 0. \quad (15c)$$

where the VEVs must satisfy the constraint (11). In section 5.2, we will express $\{v_1^2 + V_1^2, v_2^2 + V_2^2, v_3^2\}$ in terms of the parameters of the potential by using the orbital variables method.

Spectrum in the scalar neutral sector In the $H_1, H_2, H'_1, H'_2, H_3$ basis, the square mass matrix can be calculated by using $M_{ij}^2 = [\partial V(\Phi_1, \Phi_2, \Phi_3)/\partial H_i \partial H_j]|_{fields=0}$. After imposing constraints (15), we get $M_H^2 =$

$$\begin{pmatrix} 2(u_5 V_2^2 + \lambda_1 v_1^2) & (\lambda_4 - 4u_5) v_1 v_2 - 2u_5 V_1 V_2 & 2(u_5 V_1^2 + \lambda_2 v_2^2) \\ (\lambda_4 - 4u_5) v_1 v_2 - 2u_5 V_1 V_2 & 2(\lambda_1 v_1 V_1 - u_5 v_2 V_2) & (\lambda_4 - 2u_5) v_2 V_1 \\ 2(\lambda_1 v_1 V_1 - u_5 v_2 V_2) & (\lambda_4 - 2u_5) v_2 V_1 & 2(\lambda_2 v_2 V_2 - u_5 v_1 V_1) \\ \lambda_5 v_3 v_1 & & \lambda_6 v_3 v_2 \\ 2(\lambda_1 v_1 V_1 - u_5 v_2 V_2) & (\lambda_4 - 2u_5) v_2 V_1 & \lambda_5 v_3 v_1 \\ (\lambda_4 - 2u_5) v_2 V_1 & 2(\lambda_2 v_2 V_2 - u_5 v_1 V_1) & \lambda_6 v_3 v_2 \\ 2(\lambda_1 V_1^2 + u_5 v_2^2) & (\lambda_4 - 4u_5) V_1 V_2 - 2u_5 v_1 v_2 & \lambda_5 v_3 V_1 \\ (\lambda_4 - 4u_5) V_1 V_2 - 2u_5 v_1 v_2 & 2(\lambda_2 V_2^2 + u_5 v_1^2) & \lambda_6 u V_2 \\ \lambda_5 v_3 V_1 & \lambda_6 u V_2 & 2\lambda_3 v_3^2 \end{pmatrix}$$

where the term $u_5 = -(\lambda_7 + 2\lambda_{10})/4$ has been used. This mass matrix has zero determinant providing us with a Goldstone Boson G_1 and four massive scalar fields. The analytic mass values are not easy to find, but in the approximation $V_1 \sim V_2 \gg v_1 \sim v_2 \sim v_3$ they are

$$M_{he_1}^2 \approx \lambda_2 V_2^2 + \lambda_1 V_1^2 + \sqrt{(\lambda_1 V_1^2 - \lambda_2 V_2^2)^2 + (\lambda_4 + \lambda_7 + 2\lambda_{10})^2 V_1^2 V_2^2}, \quad (16)$$

$$M_{he_2}^2 \approx \lambda_2 V_2^2 + \lambda_1 V_1^2 - \sqrt{(\lambda_1 V_1^2 - \lambda_2 V_2^2)^2 + (\lambda_4 + \lambda_7 + 2\lambda_{10})^2 V_1^2 V_2^2}, \quad (17)$$

$$M_{he_3}^2 \approx -\frac{(\lambda_7 + 2\lambda_{10}) (V_2^2 + V_1^2)}{2}, \quad (18)$$

$$M_{he_4}^2 \approx 2\lambda_3 v_3^2 \quad \text{with} \quad he_4 \approx H_3, \quad (19)$$

where the scalar he_4 is light and can be identified as the SM Higgs Boson scalar. In order to have positive masses for all the former scalars, the following constraints must be satisfied

$$\lambda_1, \lambda_2, \lambda_3 > 0, \quad (20a)$$

$$4\lambda_1 \lambda_2 > (\lambda_4 + \lambda_7 + 2\lambda_{10})^2, \quad (20b)$$

$$(\lambda_7 + 2\lambda_{10}) < 0. \quad (20c)$$

Spectrum in the pseudoscalar neutral sector In the $A_1, A_2, A'_1, A'_2, A_3$ basis the square mass matrix is given by $M_A^2 =$

$$\begin{pmatrix} 2u_5 V_2^2 - \lambda_{10} v_2^2 & \lambda_{10} v_1 v_2 - 2u_5 V_1 V_2 & \frac{\lambda_7 v_2 V_2}{2} \\ \lambda_{10} v_1 v_2 - 2u_5 V_1 V_2 & 2u_5 V_1^2 - \lambda_{10} v_1^2 & -\frac{\lambda_7 v_2 V_1}{2} \\ \frac{\lambda_7 v_2 V_2}{2} & -\frac{\lambda_7 v_2 V_1}{2} & 2u_5 v_2^2 - \lambda_{10} V_2^2 \\ -\frac{\lambda_7 v_2 V_1}{2} & \frac{\lambda_7 v_1^2 V_1}{2} & \lambda_{10} V_1 V_2 - 2u_5 v_1 v_2 \\ 0 & 0 & 0 \\ -\frac{\lambda_7 v_2 V_1}{2} & 0 & 0 \\ \frac{\lambda_7 v_1^2 V_1}{2} & 0 & 0 \\ \lambda_{10} V_1 V_2 - 2u_5 v_1 v_2 & 0 & 0 \\ 2u_5 v_1^2 - \lambda_{10} V_1^2 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

which is a rank-2 matrix, giving three Goldstone Bosons and two heavy pseudoscalar particles with masses given by

$$M_{ho_1}^2 \approx -\frac{(\lambda_7 + 2\lambda_{10})(V_1^2 + V_2^2)}{2}, \quad (21)$$

$$M_{ho_2}^2 \approx -\lambda_{10}(V_1^2 + V_2^2), \quad (22)$$

where $M_{ho_1}^2 > 0$ due to the constraint (20c)], and the condition $M_{ho_2}^2 > 0$ implies the new constraint

$$\lambda_{10} < 0. \quad (23)$$

Spectrum in the charged scalar sector In the $\phi_1^\pm, \phi_2^\pm, \phi_3^\pm$ basis, the square mass matrix is given by $M_\phi^2 =$

$$\frac{1}{2} \begin{pmatrix} 4u_5 (V_2^2 + v_2^2) + \lambda_8 v_3^2 & -4u_5 (V_1 V_2 + v_1 v_2) \\ -4u_5 (V_1 V_2 + v_1 v_2) & 4u_5 (V_1^2 + v_1^2) + \lambda_9 v_3^2 \\ \lambda_8 v_3 v_1 & \lambda_9 v_3 v_2 \\ \lambda_8 v_3 V_1 & \lambda_9 v_3 V_2 \\ \lambda_8 v_3 v_1 & \lambda_8 v_3 V_1 \\ \lambda_9 v_3 v_2 & \lambda_9 v_3 V_2 \\ \lambda_9 v_2^2 + \lambda_8 v_1^2 & \lambda_9 v_2 V_2 + \lambda_8 v_1 V_1 \\ \lambda_9 v_2 V_2 + \lambda_8 v_1 V_1 & \lambda_9 V_2^2 + \lambda_8 V_1^2 \end{pmatrix} \quad (24)$$

with u_5 defined above. The mass matrix (24) is a rank-2 matrix, implying the existence of four Goldstone Bosons and four massive charged scalars, with masses given by

$$M_{h_1^\pm}^2 \approx \frac{\lambda_8 V_1^2 + \lambda_9 V_2^2}{2} > 0, \quad (25)$$

$$M_{h_2^\pm}^2 \approx -\frac{(\lambda_7 + 2\lambda_{10})(V_1^2 + V_2^2)}{2}, \quad (26)$$

where again we have $M_{h_2^\pm}^2 > 0$ due to the constraint (20c).

Counting Goldstone Bosons we have a total of eight: an scalar and three pseudoscalars which are used to provide with masses to four electrically neutral gauge bosons (Z^0, Z'^0, K^0 and \bar{K}^0), and four charged ones which are used to provide with masses to W^\pm and to K^\pm . This shows the consistency of our analysis.

3.2.2 Vacuum structure with $v_2 = V_2 = 0, v_1, V_1 \neq 0$

In this section we are going to study the other independent structure given in Table 1, where the LD between $\langle \Phi_1 \rangle$ and

$\langle \Phi_2 \rangle$ must be respected. The VEV configuration structure is

$$\langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \\ V_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad \langle \Phi_3 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_3 \\ 0 \\ 0 \end{pmatrix}. \quad (27)$$

In the tree level approximation, the constraint equations are now

$$\mu_1^2 + \lambda_1 (V_1^2 + v_1^2) + \frac{\lambda_5}{2} v_3^2 = 0, \quad (28a)$$

$$\mu_3^2 + \frac{\lambda_5}{2} (V_1^2 + v_1^2) + \lambda_3 v_3^2 = 0, \quad (28b)$$

where there is no a constraint relation for μ_2^2 , which now becomes a free parameter of the model.

Spectrum in the scalar neutral sector The first result obtained is that the fields H_1, H'_1 and H_3 do not mix with H_2 and H'_2

In the H_1, H'_1, H_3 basis, the square mass matrix is

$$M_{e_2}^2 = \begin{pmatrix} 2\lambda_1 v_1^2 & 2\lambda_1 v_1 V_1 & \lambda_5 v_1 v_3 \\ 2\lambda_1 v_1 V_1 & 2\lambda_1 V_1^2 & \lambda_5 v_3 V_1 \\ \lambda_5 v_1 v_3 & \lambda_5 v_3 V_1 & 2\lambda_3 v_3^2 \end{pmatrix}, \quad (29)$$

which is a rank-2 matrix, implying the existence of one Goldstone Boson.

Now, in the H_2, H'_2 basis, the rank-2 mass matrix is.

$$M_{e_3}^2 = \begin{pmatrix} \frac{\lambda_4 V_1^2 + S v_1^2 + \lambda_6 v_3^2}{2} + \mu_2^2 & \frac{(\lambda_7 + 2\lambda_{10}) v_1 V_1}{2} \\ \frac{(\lambda_7 + 2\lambda_{10}) v_1 V_1}{2} & \frac{S V_1^2 + \lambda_4 v_1^2 + \lambda_6 v_3^2}{2} + \mu_2^2 \end{pmatrix}, \quad (30)$$

where $S = \lambda_4 + \lambda_7 + 2\lambda_{10}$. For this vacuum structure the analytic scalar square mass values can be calculated exactly; they are

$$M_{he'_1}^2 = \lambda_1 (v_1^2 + V_1^2) + \lambda_3 v_3^2 + \sqrt{[\lambda_1 (V_1^2 + v_1^2) - \lambda_3 v_3^2]^2 + \lambda_5^2 v_3^2 (V_1^2 + v_1^2)}, \quad (31)$$

$$M_{he'_2}^2 = \lambda_1 (v_1^2 + V_1^2) + \lambda_3 v_3^2 - \sqrt{[\lambda_1 (V_1^2 + v_1^2) - \lambda_3 v_3^2]^2 + \lambda_5^2 v_3^2 (V_1^2 + v_1^2)}, \quad (32)$$

$$M_{he'_3}^2 = \frac{(\lambda_7 + \lambda_4 + 2\lambda_{10})(V_1^2 + v_1^2) + \lambda_6 v_3^2}{2} + \mu_2^2 > 0, \quad (33)$$

$$M_{he'_4}^2 = \frac{\lambda_4 (V_1^2 + v_1^2) + \lambda_6 v_3^2}{2} + \mu_2^2 > 0. \quad (34)$$

In order to have positive masses for the first two scalars, the following constraint equations must be satisfied:

$$\lambda_1, \lambda_3 > 0 \quad \text{and} \quad 4\lambda_1 \lambda_3 > \lambda_5^2. \quad (35)$$

Notice by the way that the masses for the scalar fields (31) and (32) correspond to the masses of the CP even physical fields in the economical model [12, 13, 17], and thus he'_2 can be identified as the SM Higgs boson scalar.

Spectrum in the pseudoscalar neutral sector. In this sector the fields A_1, A'_1, A_3 do not get mass entries, becoming automatically 3 odd Goldstone Bosons. Now, in the basis A_2, A'_2 the rank-2 square mass matrix M_0^2 is:

$$\begin{pmatrix} \frac{\lambda_4 V_1^2 + (\lambda_7 + \lambda_4) v_1^2 + \lambda_6 v_3^2}{2} + \mu_2^2 & \frac{\lambda_7 v_1 V_1}{2} \\ \frac{\lambda_7 v_1 V_1}{2} & \frac{(\lambda_7 + \lambda_4) V_1^2 + \lambda_4 v_1^2 + \lambda_6 v_3^2}{2} + \mu_2^2 \end{pmatrix},$$

with eigenvalues for the physical fields given by

$$M_{ho'_1}^2 = \frac{(\lambda_7 + \lambda_4)(V_1^2 + v_1^2) + \lambda_6 v_3^2}{2} + \mu_2^2 > 0, \quad (36)$$

$$M_{ho'_2}^2 = \frac{\lambda_4(V_1^2 + v_1^2) + \lambda_6 v_3^2}{2} + \mu_2^2 > 0. \quad (37)$$

Spectrum in the charged scalar sector In the $\phi_1^\pm, \phi_2^\pm, \phi_3^\pm, \phi_3^{\prime\pm}$ basis the 4×4 square mass matrix M_c^2 is

$$\begin{pmatrix} \frac{\lambda_8 v_3^2}{2} & 0 & \frac{(\lambda_8 v_1 v_3)}{2} & \frac{\lambda_8 v_3 V_1}{2} \\ 0 & \frac{\lambda_4 V_1^2 + \lambda_4 v_1^2 + (\lambda_9 + \lambda_6) v_3^2 + 2\mu_2^2}{2} & 0 & 0 \\ \frac{\lambda_8 v_1 v_3}{2} & 0 & \frac{\lambda_8 v_1^2}{2} & \frac{\lambda_8 v_1 V_1}{2} \\ \frac{\lambda_8 v_3 V_1}{2} & 0 & \frac{\lambda_8 v_1 V_1}{2} & \frac{\lambda_8 V_1^2}{2} \end{pmatrix},$$

which is a rank-2 mass matrix producing in this way four Goldstone Bosons. The remaining physical fields have square masses:

$$M_{h_1^\pm}^2 = \frac{\lambda_8(V_1^2 + v_1^2 + v_3^2)}{2}, \quad (38)$$

$$M_{h_2^\pm}^2 = \frac{\lambda_4(V_1^2 + v_1^2) + (\lambda_9 + \lambda_6)v_3^2}{2} + \mu_2^2 > 0, \quad (39)$$

where $h_2^\pm = \phi_2^\pm$. Now, for $M_{h_1^\pm}^2 > 0$ it must hold

$$\lambda_8 > 0. \quad (40)$$

Notice again that the masses of the two physical charged scalars coincide with that masses in the economical 3-3-1 model.

Counting Goldstone Bosons we get again a consistent spectrum.

In the following two sections we are going to derive bounds on the parameters of the scalar potential (10) that result from the following conditions:

- The potential $V(\Phi_1, \Phi_2, \Phi_3)$ must be stable,
- The potential must be able to break the symmetry $SU(3)_L \otimes U(1)_X$ down to $U(1)_Q$, in a consistent way.

4 Stability of the scalar Potential

The scalar potential $V(\Phi_1, \Phi_2, \Phi_3)$ in (10) is stable if it is bounded from below; this guarantees the existence of a global minimum in the potential. The stability of the scalar potential turns out to be independent of the values taken by the VEV: v_1, v_2, v_3, V_1 and V_2 , as it is going to be shown in the following analysis. In other words, the results obtained below are valid, independent of the vacuum structure chosen.

4.1 The orbital variables

The most general gauge invariant and renormalizable scalar potential $V(\Phi_1, \Phi_2, \Phi_3)$ in (10), that does not contain the cubic term, for the three Higgs scalar triplets Φ_1, Φ_2 , and Φ_3 , is an Hermitian linear combination of terms of the form

$$\Phi_i^\dagger \Phi_j, \quad (\Phi_i^\dagger \Phi_j)(\Phi_k^\dagger \Phi_l), \quad (41)$$

where $i, j, k, l \in 1, 2, 3$.

Following the method presented in appendix A, it is convenient to discuss the properties of the scalar potential, such as its stability and its spontaneous symmetry breaking, in terms of gauge invariant expressions. For this purpose we arrange the $SU(3)_L$ invariant scalar products into the the following three 2×2 hermitian matrices

$$\begin{aligned} \underline{K} &= \begin{pmatrix} \Phi_1^\dagger \Phi_1 & \Phi_2^\dagger \Phi_1 \\ \Phi_1^\dagger \Phi_2 & \Phi_2^\dagger \Phi_2 \end{pmatrix}, & \underline{L} &= \begin{pmatrix} \Phi_1^\dagger \Phi_1 & \Phi_3^\dagger \Phi_1 \\ \Phi_1^\dagger \Phi_3 & \Phi_3^\dagger \Phi_3 \end{pmatrix}, \\ \underline{M} &= \begin{pmatrix} \Phi_2^\dagger \Phi_2 & \Phi_3^\dagger \Phi_2 \\ \Phi_2^\dagger \Phi_3 & \Phi_3^\dagger \Phi_3 \end{pmatrix}, \end{aligned} \quad (42)$$

where according with Eq. (125), each matrix is related to the following four real parameters

$$\underline{K}: \begin{cases} \Phi_1^\dagger \Phi_1 = (K_0 + K_3)/2, & \Phi_2^\dagger \Phi_2 = (K_0 - K_3)/2, \\ \Phi_1^\dagger \Phi_2 = (K_1 + i K_2)/2, & \Phi_2^\dagger \Phi_1 = (K_1 - i K_2)/2, \end{cases} \quad (43a)$$

$$\underline{L}: \begin{cases} \Phi_1^\dagger \Phi_1 = (L_0 + L_3)/2, & \Phi_3^\dagger \Phi_3 = (L_0 - L_3)/2, \\ \Phi_1^\dagger \Phi_3 = (L_1 + i L_2)/2, & \Phi_3^\dagger \Phi_1 = (L_1 - i L_2)/2, \end{cases} \quad (43b)$$

$$\underline{M}: \begin{cases} \Phi_2^\dagger \Phi_2 = (M_0 + M_3)/2, & \Phi_3^\dagger \Phi_3 = (M_0 - M_3)/2, \\ \Phi_2^\dagger \Phi_3 = (M_1 + i M_2)/2, & \Phi_3^\dagger \Phi_2 = (M_1 - i M_2)/2, \end{cases} \quad (43c)$$

with the constraints

$$K_0 \geq 0, \quad K_0^2 - K_1^2 - K_2^2 - K_3^2 = K_0^2 - \mathbf{K}^2 \geq 0, \quad (44a)$$

$$L_0 \geq 0, \quad L_0^2 - L_1^2 - L_2^2 - L_3^2 = L_0^2 - \mathbf{L}^2 \geq 0, \quad (44b)$$

$$M_0 \geq 0, \quad M_0^2 - M_1^2 - M_2^2 - M_3^2 = M_0^2 - \mathbf{M}^2 \geq 0. \quad (44c)$$

The scalar products $\Phi_1^\dagger \Phi_1, \Phi_2^\dagger \Phi_2$ and $\Phi_3^\dagger \Phi_3$, present in the expressions (43a), (43b) and (43c), allow us to eliminate three of the 12 variables due to the fact that

$$K_3 = L_0 - M_0, \quad (45a)$$

$$L_3 = K_0 - M_0, \quad (45b)$$

$$M_3 = K_0 - L_0, \quad (45c)$$

ending up with only the following nine real orbital variables, used to describe the full scalar potential

$$K_0, L_0, M_0, K_1, K_2, L_1, L_2, M_1, M_2. \quad (46)$$

With the help of the former variables, the scalar potential (10) may be written as

$$V(\Phi_1, \Phi_2, \Phi_3) = (V_{2K} + V_{4K}) + (V_{2L} + V_{4L}) + (V_{2M} + V_{4M}), \quad (47)$$

where as can be seen, the general space splits as the direct sum of three subspaces, due to the particular simple form of the scalar potential in (10) and to the fact that $\langle \Phi_3 \rangle$ is

orthogonal to $\langle \Phi_1 \rangle$ and to $\langle \Phi_2 \rangle$, something which guarantees the validity of the generalized Schwartz's Inequality

$$\langle \Phi_1 | \Phi_1 \rangle \langle \Phi_2 | \Phi_2 \rangle \langle \Phi_3 | \Phi_3 \rangle \geq \langle \Phi_1 | \Phi_2 \rangle \langle \Phi_1 | \Phi_3 \rangle \langle \Phi_2 | \Phi_3 \rangle.$$

With the use of the real parameters $\xi_{k(l,m)0}$, $\xi_{k(l,m)a}$, $\eta_{k(l,m)00}$, $\eta_{k(l,m)a}$ and $\eta_{k(l,m)ab} = \eta_{k(l,m)ba}$, the following functions defined in the domain $|\mathbf{k}|, |\mathbf{l}|, |\mathbf{m}| \leq 1$.

$$J_{k2}(\mathbf{k}) = \xi_{k0} + \boldsymbol{\xi}_k \cdot \mathbf{k}, \quad (48a)$$

$$J_{k4}(\mathbf{k}) = \eta_{k00} + 2\boldsymbol{\eta}_k \cdot \mathbf{k} + \mathbf{k} \cdot E_k \cdot \mathbf{k}, \quad (48b)$$

$$J_{l2}(\mathbf{l}) = \xi_{l0} + \boldsymbol{\xi}_l \cdot \mathbf{l}, \quad (48c)$$

$$J_{l4}(\mathbf{l}) = \eta_{l00} + 2\boldsymbol{\eta}_l \cdot \mathbf{l} + \mathbf{l} \cdot E_l \cdot \mathbf{l}, \quad (48d)$$

$$J_{m2}(\mathbf{m}) = \xi_{m0} + \boldsymbol{\xi}_m \cdot \mathbf{m}, \quad (48e)$$

$$J_{m4}(\mathbf{m}) = \eta_{m00} + 2\boldsymbol{\eta}_m \cdot \mathbf{m} + \mathbf{m} \cdot E_m \cdot \mathbf{m}, \quad (48f)$$

where in according to (44)

$$\mathbf{k} = \mathbf{K}/K_0, \quad (|\mathbf{k}| \leq 1); \quad (49a)$$

$$\mathbf{l} = \mathbf{L}/L_0, \quad (|\mathbf{l}| \leq 1); \quad (49b)$$

$$\mathbf{m} = \mathbf{M}/M_0, \quad (|\mathbf{m}| \leq 1), \quad (49c)$$

for $K_0, L_0, M_0 > 0$, allows us to write the terms of potential (47).

$$V_{2K} = \xi_{k0}K_0 + \xi_{ka}K_a = K_0J_{k2}(\mathbf{k}), \quad (50a)$$

$$V_{4K} = \eta_{k00}K_0^2 + 2K_0\eta_{ka}K_a + K_a\eta_{kab}K_b = K_0^2J_{k4}(\mathbf{k}), \quad (50b)$$

$$V_{2L} = \xi_{l0}L_0 + \xi_{la}L_a = L_0J_{l2}(\mathbf{l}), \quad (50c)$$

$$V_{4L} = \eta_{l00}L_0^2 + 2L_0\eta_{la}L_a + L_a\eta_{lab}L_b = L_0^2J_{l4}(\mathbf{l}), \quad (50d)$$

$$V_{2M} = \xi_{m0}M_0 + \xi_{ma}M_a = M_0J_{m2}(\mathbf{m}), \quad (50e)$$

$$V_{4M} = \eta_{m00}M_0^2 + 2M_0\eta_{ma}M_a + M_a\eta_{mab}M_b = M_0^2J_{m4}(\mathbf{m}), \quad (50f)$$

where sum over the indices a and b from 1 to 3 must be understood. In the former expressions, the following notation has been used: $E_k = \eta_{kab}$, $E_l = \eta_{lab}$, $E_m = \eta_{mab}$. The parametrization employed in Eqs. (42)-(46) should not invalidate the stability conditions (in the strong sense) as far as sufficient conditions are concerned (necessary and sufficient conditions should be affected).

On the other hand, the parameters given in (46) does not imply that the matrix arrangements (42) can be established. In that way, the parameters (46) may help in the procedure to find the stationary points in the scalar potential, but it is necessary to verify at the end, if all matrices (42) are consistent with the stationary points found. That is the analysis given below.

4.2 Stability conditions

For the potential to be stable, it must be bounded from below. The stability is determined by the behavior of V in the limit $K_0 \rightarrow \infty$, $L_0 \rightarrow \infty$ and/or $M_0 \rightarrow \infty$; hence, by

the signs of $J_{k(l,m)2}(\mathbf{k}, \mathbf{l}, \mathbf{m})$ and $J_{k(l,m)4}(\mathbf{k}, \mathbf{l}, \mathbf{m})$ in (50), (approach which conduces only to sufficiency conditions but not to necessary conditions).

In the strong sense, the stability of the potential is guaranteed when $V \rightarrow \infty$ for \mathbf{k}, \mathbf{l} and \mathbf{m} taking any value, which means that

$$J_{k4}(\mathbf{k}), J_{l4}(\mathbf{l}), J_{m4}(\mathbf{m}) > 0 \text{ for all } |\mathbf{k}|, |\mathbf{l}|, |\mathbf{m}| \leq 1. \quad (51)$$

To assure the existence of a positive (semi-)definite value for $J_{k(l,m)4}(\mathbf{k}, \mathbf{l}, \mathbf{m})$, it is sufficient to consider its value for all the stationary points of $J_{k(l,m)4}(\mathbf{k}, \mathbf{l}, \mathbf{m})$ in the domain $|\mathbf{k}|, |\mathbf{l}|, |\mathbf{m}| < 1$, and for all stationary points on the boundary $|\mathbf{k}|, |\mathbf{l}|, |\mathbf{m}| = 1$. This holds, because the global minimum of the continuous function $J_{k(l,m)4}(\mathbf{k}, \mathbf{l}, \mathbf{m})$ is reached on the compact domain $|\mathbf{k}|, |\mathbf{l}|, |\mathbf{m}| \leq 1$, and it is located among those stationary points. This leads to bounds on $\eta_{k(l,m)00}$, $\eta_{k(l,m)a}$ and $\eta_{k(l,m)ab}$, which parametrise the quartic term $V_{4K(L,M)}$ of the potential. A detailed analysis of the stability criteria for a scalar potential can be found in Refs [3,13].

With the help of Eqs. (43) and (45), the parameters defined in (50), for the scalar potential in (10) are:

$$\begin{aligned} \xi_{k0} &= (\mu_1^2 + \mu_2^2 - \mu_3^2)/2, \quad \eta_{k00} = (\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4)/4, \\ \boldsymbol{\xi}_k &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad \boldsymbol{\eta}_k = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \\ E_k &= \begin{pmatrix} (\lambda_7 + 2\lambda_{10})/4 & 0 & 0 \\ 0 & \lambda_7/4 & 0 \\ 0 & 0 & (\lambda_1 + \lambda_2 - \lambda_3 - \lambda_4)/4 \end{pmatrix}, \\ \xi_{l0} &= (\mu_1^2 - \mu_2^2 + \mu_3^2)/2, \quad \eta_{l00} = (\lambda_1 - \lambda_2 + \lambda_3 + \lambda_5)/4, \\ \boldsymbol{\xi}_l &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad \boldsymbol{\eta}_l = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \\ E_l &= \begin{pmatrix} \lambda_8/4 & 0 & 0 \\ 0 & \lambda_8/4 & 0 \\ 0 & 0 & (\lambda_1 - \lambda_2 + \lambda_3 - \lambda_5)/4 \end{pmatrix}, \\ \xi_{m0} &= (-\mu_1^2 + \mu_2^2 + \mu_3^2)/2, \quad \eta_{m00} = (-\lambda_1 + \lambda_2 + \lambda_3 + \lambda_6)/4, \\ \boldsymbol{\xi}_m &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad \boldsymbol{\eta}_m = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \\ E_m &= \begin{pmatrix} \lambda_9/4 & 0 & 0 \\ 0 & \lambda_9/4 & 0 \\ 0 & 0 & (-\lambda_1 + \lambda_2 + \lambda_3 - \lambda_6)/4 \end{pmatrix}. \end{aligned} \quad (52)$$

The stability criteria established in the previous section allow us to bound the parameters of the potential in the following way:

For K :

$$\lambda_1 + \lambda_2 - \lambda_3 > 0, \quad (53a)$$

$$\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 > 0, \quad (53b)$$

$$\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 + \lambda_7 > 0, \quad (53c)$$

$$\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 + \lambda_7 + 2\lambda_{10} > 0, \quad (53d)$$

For L :

$$\lambda_1 - \lambda_2 + \lambda_3 > 0, \quad (54a)$$

$$\lambda_1 - \lambda_2 + \lambda_3 + \lambda_5 > 0, \quad (54b)$$

$$\lambda_1 - \lambda_2 + \lambda_3 + \lambda_5 + \lambda_8 > 0, \quad (54c)$$

The domain of orbital variables, Eqs. (44), can be written in the following form.

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} \geq 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} \geq 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} \geq 0, K_0 \geq 0, L_0 \geq 0, M_0 \geq 0, \quad (69)$$

with

$$\begin{aligned} \tilde{\mathbf{g}}_1 &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \\ \tilde{\mathbf{g}}_2 &= \begin{pmatrix} -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (70) \\ \tilde{\mathbf{g}}_3 &= \begin{pmatrix} -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}. \end{aligned}$$

The trivial configuration $\tilde{\mathbf{P}} = 0$ is a stationary point of the potential with $V = 0$, as a direct consequence of the definitions.

For the discussion of the stationary points of V , we must distinguish among the following cases

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} > 0; \quad (71a)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} > 0; \quad (71b)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} > 0; \quad (71c)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} = 0; \quad (71d)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} > 0; \quad (71e)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} = 0; \quad (71f)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} > 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} = 0; \quad (71g)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} = 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} = 0. \quad (71h)$$

The stationary points of V in the inner part of the domain, cases (71a), (71c), (71d) and (71g), imply linear independence between $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$, something which is not allowed. The global minimum for the case (71h) implies LD among $\langle \Phi_1 \rangle$, $\langle \Phi_2 \rangle$ and $\langle \Phi_3 \rangle$, not allowed either. Case (71e) implies $\langle \Phi_1 \rangle = 0$. So, the only two cases of concern for us here are (71b) corresponding to the general vacuum structure with $v_1, v_2, v_3, V_1, V_2 \neq 0$ studied

in Sect. (3.2.1), and case (71f) which corresponds to the vacuum structure $\langle \Phi_2 \rangle = 0$ studied in Sect. (3.2.2).

In general, the stationary points of the scalar potential in (68), for any domain in (71), are stationary points of the function

$$\tilde{F}(\tilde{\mathbf{P}}, u, v, w) = V - u\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} - v\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} - w\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}}, \quad (72)$$

where u, v and w are Lagrange multipliers. The relevant stationary points of \tilde{F} are thus given as solutions to the equation

$$(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)\tilde{\mathbf{P}} = -\frac{1}{2}\tilde{\xi}; \quad \text{with} \quad (73a)$$

$$\tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{P}} \geq 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{P}} \geq 0, \tilde{\mathbf{P}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{P}} \geq 0, \quad (73b)$$

and

$$K_0, L_0, M_0 > 0, \quad (74)$$

with the inequality (> 0) in Eq. (73b) taking place, for the case when the Lagrange multipliers are excluded. For regular values of u, v and w , with the determinant

$$\det(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3) \neq 0$$

we have

$$\tilde{\mathbf{P}} = -\frac{1}{2}(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\xi}. \quad (75)$$

The Lagrange multipliers are thus obtained by inserting (75) in the constraint Eqs. (73b):

$$\begin{aligned} \tilde{\xi}(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\mathbf{g}}_1(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\xi} &= 0, \\ \tilde{\xi}(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\mathbf{g}}_2(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\xi} &= 0, \\ \tilde{\xi}(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\mathbf{g}}_3(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3)^{-1}\tilde{\xi} &= 0, \\ \text{and } K_0, L_0, M_0 > 0. \end{aligned} \quad (76)$$

Additionally, there may be up to 9 values $u = \tilde{\mu}_a$ (and also 9 values for $v = \tilde{\mu}_a$, and for $w = \tilde{\mu}_a$) with $a = 1, \dots, 9$, for which $\det(\tilde{E} - u\tilde{\mathbf{g}}_1 - v\tilde{\mathbf{g}}_2 - w\tilde{\mathbf{g}}_3) = 0$. Depending on the form of the potential; some, or all of them, may lead to exceptional solutions of (73a).

For *any* stationary point of the potential we have

$$V|_{stat} = \frac{1}{2}\tilde{\mathbf{P}} \cdot \tilde{\xi} = -\tilde{\mathbf{P}} \cdot \tilde{E} \cdot \tilde{\mathbf{P}}. \quad (77)$$

Suppose now that the strong stability condition (51) holds. Then (77) gives for non-trivial stationary points where $\tilde{\mathbf{P}} \neq 0$:

$$V|_{stat} < 0. \quad (78)$$

Firstly, in the context that only one Lagrange multiplier is non-zero, for instance $u_p \neq 0, v = w = 0$ in (73a), let's consider $\tilde{\mathbf{p}} = (p_{k_0}, p_{l_0}, p_{m_0}, p_{k_1}, p_{k_2}, p_{l_1}, p_{l_2}, p_{m_1}, p_{m_2})^T$ be a stationary point for this case. Then, from (68) and (73a)

we have

$$\left. \frac{\partial V}{\partial K_0} \right|_{\substack{\tilde{\mathbf{P}}_{\{1\}} \text{ fixed,} \\ \tilde{\mathbf{P}} = \tilde{\mathbf{p}}}} = 2 u_p p_{k_0}, \quad (79a)$$

$$\left. \frac{\partial V}{\partial L_0} \right|_{\substack{\tilde{\mathbf{P}}_{\{2\}} \text{ fixed,} \\ \tilde{\mathbf{P}} = \tilde{\mathbf{p}}}} = 2 u_p p_{l_0}, \quad (79b)$$

$$\left. \frac{\partial V}{\partial M_0} \right|_{\substack{\tilde{\mathbf{P}}_{\{3\}} \text{ fixed,} \\ \tilde{\mathbf{P}} = \tilde{\mathbf{p}}}} = 2 u_p p_{m_0}, \quad (79c)$$

where the notation established in (67) has been used. For the analysis which follows, only the most convenient partial derivative from (79) is chosen, in such a way that only two, out of the three values in (45) lower down, and also that the inequalities (44) hold for the new points; (for example, let us take M_0 . From (45) we have $(K_3, L_3) \rightarrow (K'_3 = K_3 - \Delta M_0), (L'_3 = L_3 - \Delta M_0)$. Where $|K'_3| < |K_3|$ and $|L'_3| < |L_3|$, if $K_3, L_3 > 0$). If $u_p < 0$, there are points $\tilde{\mathbf{P}}$ with $K_0 > p_{k_0}$ (or $L_0 > p_{l_0}, M_0 > p_{m_0}$), $\tilde{\mathbf{P}}_{\{1\}} = \tilde{\mathbf{p}}_{\{1\}}$ (or $\tilde{\mathbf{P}}_{\{2\}} = \tilde{\mathbf{p}}_{\{2\}}, \tilde{\mathbf{P}}_{\{3\}} = \tilde{\mathbf{p}}_{\{3\}}$) for which the potential decreases in its neighborhood, and as a consequence cannot be a minimum. We conclude that in a theory with the required EWSB, a stationary point coming from an unique no null Lagrange multiplier (for example $u_0 \neq 0, v = w = 0$), to be a global minimum candidate, it must hold

$$u_0 > 0. \quad (80)$$

Secondly, for the stationary points $\tilde{\mathbf{p}}$ and $\tilde{\mathbf{q}}$, we have in general, from (73a) and (77), the following relation

$$\begin{aligned} V(\tilde{\mathbf{p}}) - V(\tilde{\mathbf{q}}) &= \frac{1}{2} \tilde{\mathbf{p}} \cdot \tilde{\boldsymbol{\xi}} - \frac{1}{2} \tilde{\mathbf{q}} \cdot \tilde{\boldsymbol{\xi}} \\ &= \tilde{\mathbf{p}} \cdot (u_q \tilde{\mathbf{g}}_1 + v_q \tilde{\mathbf{g}}_2 + w_q \tilde{\mathbf{g}}_3 - \tilde{\mathbf{E}}) \cdot \tilde{\mathbf{q}} \\ &\quad - \tilde{\mathbf{q}} \cdot (u_p \tilde{\mathbf{g}}_1 + v_p \tilde{\mathbf{g}}_2 + w_p \tilde{\mathbf{g}}_3 - \tilde{\mathbf{E}}) \cdot \tilde{\mathbf{p}} \\ &= (u_q - u_p) \tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{q}} + (v_q - v_p) \tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{q}} \\ &\quad + (w_q - w_p) \tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{q}}, \end{aligned} \quad (81)$$

where $\tilde{\mathbf{p}}$ and $\tilde{\mathbf{q}}$ are vectors on the forward light cone, and $\tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_1 \cdot \tilde{\mathbf{q}}, \tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_2 \cdot \tilde{\mathbf{q}}, \tilde{\mathbf{p}} \cdot \tilde{\mathbf{g}}_3 \cdot \tilde{\mathbf{q}}$ are always non-negative.

In Table 2, an exhaustive of all the possible stationary points of the potential are presented (even if it includes unphysical VEVs). Lagrange multipliers coming from solutions of Eq. (73a) belonging to regulars and exceptional values for the cases are stated. Those cases of Lagrange multipliers giving the same stationary point, only one of them were included in Table 2. Solutions which imply specific relations among the parameters $\mu_1^2, \mu_2^2, \mu_3^2$ were excluded too (see for example [13]). Incidentally, it is important to say that all stationary points written in Table 2 give physical VEVs, i.e, they are consistent in relation with matrices (42).

To end our analysis, let us apply our findings to the two independent vacuum structures given by the constraints (71b) and (71f), which were studied in detail in Sects. (3.2.1) and (3.2.2) respectively.

5.1 Case $v_2 = V_2 = 0$

We want a global minimum with the configuration (27), which implies solutions satisfying (71f). For this purpose we use the results in Table (2), the stability conditions stated in Sec. (4.2) and taking into account expression (81). In the case of an unique no null Lagrange multiplier, the restriction in (80) can be used. In that way, the conditions found below are sufficient (but not necessary) to have the minimum of the scalar potential (10) at $\tilde{\mathbf{P}}_1$.

From the Table 2, we want the global minimum be associated to Lagrange multiplier u_1 (where w_2 gives the same stationary point) which does not coincide with solutions inside the forward light cone (71a).

Let's assume the following conditions

$$\lambda_4 > 0, \quad \lambda_5 < 0 \quad \text{and} \quad \lambda_6 > 0, \quad (82)$$

which in combination with the inequalities (56), (57) and (65), we have that $\lambda_5 \mu_3^2 - 2\lambda_3 \mu_1^2 > 0$, $\lambda_5 \mu_1^2 - 2\lambda_1 \mu_3^2 > 0$, and $4\lambda_1 \lambda_3 - \lambda_5^2 > 0$, hence $K_{01} > 0, M_{01} > 0$ and $L_{01} > 0$. Additionally, with same arguments, we can verify that the Lagrange multipliers $v_{13}, v_{15} < 0$.

As you can see, there is not inconvenient to impose the following condition

$$u_1 > 0, \quad (83)$$

as is required by the global minimum condition (80). Therefore, for the moment, the point $\tilde{\mathbf{P}}_1$ satisfy all requirements to be a stationary point. The other aspect to take into account, it is to show that this point is the global minimum of potential. For that, let's see the other Lagrange multipliers and their points, and to establish conditions over them such that the global are not found there.

For example, if we assume

$$\lambda_7 + 2\lambda_{10} > 0, \quad \lambda_7 > 0, \quad \lambda_8 > 0 \quad \lambda_9 > 0, \quad (84)$$

it immediately discards out the points $\tilde{\mathbf{P}}_5, \tilde{\mathbf{P}}_6, \tilde{\mathbf{P}}_7$ and $\tilde{\mathbf{P}}_8$ as minimal global points of potential, because the corresponding Lagrange multipliers u_5, u_6, v_7 and w_8 are negative numbers.

If we assume

$$\lambda_6 \mu_2^2 - 2\lambda_2 \mu_3^2 > 0, \quad (85)$$

$$\lambda_6 \mu_3^2 - 2\lambda_3 \mu_2^2 < 0, \quad (86)$$

it gives either the condition $K_{03} > 0$ or $L_{03} > 0$, but not both conditions satisfied simultaneously. And, in similar way, considering the case

$$\lambda_4 \mu_2^2 - 2\lambda_2 \mu_1^2 > 0, \quad (87)$$

$$\lambda_4 \mu_1^2 - 2\lambda_1 \mu_2^2 < 0 \quad (88)$$

we conclude that L_4, M_4 are not positive numbers simultaneously. Then, $\tilde{\mathbf{P}}_3$ and $\tilde{\mathbf{P}}_4$ are not stationary points of potential.

From (65), (87) and (85) we see that the Lagrange multipliers $u_{14}, w_{14} < 0$. Thus, the minimum of potential is not present at $\tilde{\mathbf{P}}_{14}$.

Domains	Lagrange Multipliers	Stationary Points
$\tilde{P}_1 \cdot \tilde{g}_1 \cdot \tilde{P}_1 = 0,$ $\tilde{P}_1 \cdot \tilde{g}_2 \cdot \tilde{P}_1 = 4 K_{01} M_{01},$ $\tilde{P}_1 \cdot \tilde{g}_3 \cdot \tilde{P}_1 = 0.$	$u_1 = \frac{\mu_2^2 + \lambda_4 K_{01} + \lambda_6 M_{01}}{4 K_{01}}, \quad w_1 = 0$	$\underline{\tilde{P}}_1 = \begin{pmatrix} K_{01} = \frac{\lambda_5 \mu_3^2 - 2 \lambda_3 \mu_1^2}{4 \lambda_1 \lambda_3 - \lambda_5^2} \\ L_{01} = K_{01} + M_{01} \\ M_{01} = \frac{\lambda_5 \mu_1^2 - 2 \lambda_1 \mu_3^2}{4 \lambda_1 \lambda_3 - \lambda_5^2} \end{pmatrix}$
	$u_2 = 0, \quad w_2 = \frac{\mu_2^2 + \lambda_4 K_{01} + \lambda_6 M_{01}}{4 M_{01}}$	$\underline{\tilde{P}}_2 = \tilde{P}_1$
$\tilde{P}_3 \cdot \tilde{g}_1 \cdot \tilde{P}_3 = 0,$ $\tilde{P}_3 \cdot \tilde{g}_2 \cdot \tilde{P}_3 = 0,$ $\tilde{P}_3 \cdot \tilde{g}_3 \cdot \tilde{P}_3 = 4 K_{03} L_{03}.$	$u_3 = \frac{\mu_1^2 + \lambda_4 K_{03} + \lambda_5 L_{03}}{4 K_{03}}, \quad v_3 = 0$	$\underline{\tilde{P}}_3 = \begin{pmatrix} K_{03} = \frac{\lambda_6 \mu_3^2 - 2 \lambda_3 \mu_2^2}{4 \lambda_2 \lambda_3 - \lambda_6^2} \\ L_{03} = \frac{\lambda_6 \mu_2^2 - 2 \lambda_2 \mu_3^2}{4 \lambda_2 \lambda_3 - \lambda_6^2} \\ K_{03} + L_{03} \end{pmatrix}$
$\tilde{P}_4 \cdot \tilde{g}_1 \cdot \tilde{P}_4 = 4 L_4 M_4,$ $\tilde{P}_4 \cdot \tilde{g}_2 \cdot \tilde{P}_4 = 0,$ $\tilde{P}_4 \cdot \tilde{g}_3 \cdot \tilde{P}_4 = 0.$	$v_4 = 0, \quad w_4 = \frac{\mu_3^2 + \lambda_5 L_4 + \lambda_6 M_4}{4 M_4}$	$\underline{\tilde{P}}_4 = \begin{pmatrix} L_4 + M_4 \\ L_4 = \frac{\lambda_4 \mu_2^2 - 2 \lambda_2 \mu_1^2}{4 \lambda_1 \lambda_2 - \lambda_4^2} \\ M_4 = \frac{\lambda_4 \mu_1^2 - 2 \lambda_1 \mu_2^2}{4 \lambda_1 \lambda_2 - \lambda_4^2} \end{pmatrix}$
$\tilde{P}_{\{5,6\}} \cdot \tilde{g}_1 \cdot \tilde{P}_{\{5,6\}} = 0,$ $\tilde{P}_{\{5,6\}} \cdot \tilde{g}_2 \cdot \tilde{P}_{\{5,6\}} \neq 0,$ $\tilde{P}_{\{5,6\}} \cdot \tilde{g}_3 \cdot \tilde{P}_{\{5,6\}} \neq 0.$	Exceptional solution: $u_5 = -\frac{\lambda_7 + 2 \lambda_{10}}{4}$	$\underline{\tilde{P}}_5$
	Exceptional solution: $u_6 = -\frac{\lambda_7}{4}$	$\underline{\tilde{P}}_6$
$\tilde{P}_7 \cdot \tilde{g}_1 \cdot \tilde{P}_7 \neq 0,$ $\tilde{P}_7 \cdot \tilde{g}_2 \cdot \tilde{P}_7 = 0,$ $\tilde{P}_7 \cdot \tilde{g}_3 \cdot \tilde{P}_7 \neq 0.$	Exceptional solution: $v_7 = -\frac{\lambda_8}{4}$	$\underline{\tilde{P}}_7$
$\tilde{P}_8 \cdot \tilde{g}_1 \cdot \tilde{P}_8 \neq 0,$ $\tilde{P}_8 \cdot \tilde{g}_2 \cdot \tilde{P}_8 \neq 0,$ $\tilde{P}_8 \cdot \tilde{g}_3 \cdot \tilde{P}_8 = 0.$	Exceptional solution: $w_8 = -\frac{\lambda_9}{4}$	$\underline{\tilde{P}}_8$
$\tilde{P}_9 \cdot \tilde{g}_1 \cdot \tilde{P}_9 = 0,$ $\tilde{P}_9 \cdot \tilde{g}_2 \cdot \tilde{P}_9 = 0,$ $\tilde{P}_9 \cdot \tilde{g}_3 \cdot \tilde{P}_9 = 0.$	Exceptional solution: $u_9 = -\frac{\lambda_7 + 2 \lambda_{10}}{4},$ $v_9 = \frac{\mu_3^2 + \lambda_5 L_9 + \lambda_6 M_9}{4 L_9}, \quad w_9 = 0$	$\underline{\tilde{P}}_9 = \begin{pmatrix} L_9 + M_9 \\ L_9 = \frac{(\lambda_4 + \lambda_7 + 2 \lambda_{10}) \mu_2^2 - 2 \lambda_2 \mu_1^2}{4 \lambda_1 \lambda_2 - (\lambda_4 + \lambda_7 + 2 \lambda_{10})^2} \\ M_9 = \frac{(\lambda_4 + \lambda_7 + 2 \lambda_{10}) \mu_1^2 - 2 \lambda_1 \mu_2^2}{4 \lambda_1 \lambda_2 - (\lambda_4 + \lambda_7 + 2 \lambda_{10})^2} \\ \pm 2 \sqrt{L_9 M_9} \end{pmatrix}$
$\tilde{P}_{10} \cdot \tilde{g}_1 \cdot \tilde{P}_{10} = 0,$ $\tilde{P}_{10} \cdot \tilde{g}_2 \cdot \tilde{P}_{10} = 0,$ $\tilde{P}_{10} \cdot \tilde{g}_3 \cdot \tilde{P}_{10} = 0.$	Exceptional solution: $u_{10} = -\frac{\lambda_7}{4},$ $v_{10} = \frac{\mu_3^2 + \lambda_5 L_{10} + \lambda_6 M_{10}}{4 L_{10}}, \quad w_{10} = 0$	$\underline{\tilde{P}}_{10} = \begin{pmatrix} L_{10} + M_{10} \\ L_{10} = \frac{(\lambda_4 + \lambda_7) \mu_2^2 - 2 \lambda_2 \mu_1^2}{4 \lambda_1 \lambda_2 - (\lambda_4 + \lambda_7)^2} \\ M_{10} = \frac{(\lambda_4 + \lambda_7) \mu_1^2 - 2 \lambda_1 \mu_2^2}{4 \lambda_1 \lambda_2 - (\lambda_4 + \lambda_7)^2} \\ 0 \\ \pm 2 \sqrt{L_{10} M_{10}} \end{pmatrix}$
$\tilde{P}_{11} \cdot \tilde{g}_1 \cdot \tilde{P}_{11} = 0,$ $\tilde{P}_{11} \cdot \tilde{g}_2 \cdot \tilde{P}_{11} = 0,$ $\tilde{P}_{11} \cdot \tilde{g}_3 \cdot \tilde{P}_{11} = 0.$	Exceptional solution: $u_{11} = \frac{\mu_2^2 + \lambda_4 K_{11} + \lambda_6 M_{11}}{4 K_{11}},$ $v_{11} = -\frac{\lambda_8}{4}, \quad w_{11} = 0$	$\underline{\tilde{P}}_{11} = \begin{pmatrix} K_{11} = \frac{(\lambda_5 + \lambda_8) \mu_3^2 - 2 \lambda_3 \mu_1^2}{4 \lambda_1 \lambda_3 - (\lambda_5 + \lambda_8)^2} \\ K_{11} + M_{11} \\ M_{11} = \frac{(\lambda_5 + \lambda_8) \mu_1^2 - 2 \lambda_1 \mu_3^2}{4 \lambda_1 \lambda_3 - (\lambda_5 + \lambda_8)^2} \\ 0 \\ \pm 2 \sqrt{K_{11} M_{11}} \end{pmatrix}$
$\tilde{P}_{12} \cdot \tilde{g}_1 \cdot \tilde{P}_{12} = 0,$ $\tilde{P}_{12} \cdot \tilde{g}_2 \cdot \tilde{P}_{12} = 0,$ $\tilde{P}_{12} \cdot \tilde{g}_3 \cdot \tilde{P}_{12} = 0.$	Exceptional solution: $u_{12} = \frac{\mu_1^2 + \lambda_4 K_{12} + \lambda_5 L_{12}}{4 K_{12}},$ $v_{12} = -\frac{\lambda_9}{4}, \quad w_{12} = 0$	$\underline{\tilde{P}}_{12} = \begin{pmatrix} K_{12} = \frac{(\lambda_6 + \lambda_9) \mu_3^2 - 2 \lambda_3 \mu_2^2}{4 \lambda_2 \lambda_3 - (\lambda_6 + \lambda_9)^2} \\ L_{12} = \frac{(\lambda_6 + \lambda_9) \mu_2^2 - 2 \lambda_2 \mu_3^2}{4 \lambda_2 \lambda_3 - (\lambda_6 + \lambda_9)^2} \\ K_{12} + L_{12} \\ 0 \\ 0 \\ 0 \\ \pm 2 \sqrt{K_{12} L_{12}} \end{pmatrix}$
$\tilde{P} \cdot \tilde{g}_1 \cdot \tilde{P} = 0,$ $\tilde{P} \cdot \tilde{g}_2 \cdot \tilde{P} = 0,$ $\tilde{P} \cdot \tilde{g}_3 \cdot \tilde{P} = 0.$	$u_{13} = \frac{\lambda_4 \mu_1^2 - 2 \lambda_1 \mu_2^2}{4 \mu_1^2}, \quad v_{13} = \frac{\lambda_5 \mu_1^2 - 2 \lambda_1 \mu_3^2}{4 \mu_1^2}, \quad w_{13} = 0$	$\underline{\tilde{P}}_{13} = \begin{pmatrix} -\frac{\mu_1^2}{2 \lambda_1} \\ -\frac{\mu_1^2}{2 \lambda_1} \\ 0 \end{pmatrix}$
	$u_{14} = \frac{\lambda_4 \mu_2^2 - 2 \lambda_2 \mu_1^2}{4 \mu_2^2}, \quad v_{14} = 0, \quad w_{14} = \frac{\lambda_6 \mu_2^2 - 2 \lambda_2 \mu_3^2}{4 \mu_2^2}$	$\underline{\tilde{P}}_{14} = \begin{pmatrix} -\frac{\mu_2^2}{2 \lambda_2} \\ 0 \\ -\frac{\mu_2^2}{2 \lambda_2} \end{pmatrix}$
	$u_{15} = 0, \quad v_{15} = \frac{\lambda_5 \mu_3^2 - 2 \lambda_3 \mu_1^2}{4 \mu_3^2}, \quad w_{15} = \frac{\lambda_6 \mu_3^2 - 2 \lambda_3 \mu_2^2}{4 \mu_3^2}$	$\underline{\tilde{P}}_{15} = \begin{pmatrix} 0 \\ -\frac{\mu_3^2}{2 \lambda_3} \\ -\frac{\mu_3^2}{2 \lambda_3} \end{pmatrix}$

Table 2. Lagrange Multipliers. The stationary points ($\underline{\tilde{P}}$) were underlined indicating that only the upper non-zero entries of column vector are written explicitly, with the remaining ones entries filled by zeros.

We derived above that $v_{13} < 0$ and $v_{15} < 0$, that together with conditions (65) and (82), it is easy to verify that $u_1 - u_{13} = [v_{13}(\lambda_6 \mu_1^2 - \lambda_5 \mu_2^2)] / (4\mu_3^2 v_{15}) > 0$, which implies that $u_1 > u_{13}$. In the same way, $w_2 - w_{15} = [v_{15}(\lambda_4 \mu_3^2 - \lambda_5 \mu_2^2)] / (4\mu_1^2 v_{13}) > 0$, that is, $w_2 > w_{15}$. Therefore, in the points $\tilde{\mathbf{P}}_{13}$ and $\tilde{\mathbf{P}}_{15}$ the global minimum are not found.

Remain to see the points $\tilde{\mathbf{P}}_9, \tilde{\mathbf{P}}_{10}, \tilde{\mathbf{P}}_{11}$ and $\tilde{\mathbf{P}}_{12}$. In order to discard these points as global minima, we can proceed in the same way as we did with the points $\tilde{\mathbf{P}}_3$ and $\tilde{\mathbf{P}}_4$. Let's consider the numerator of $L_9, L_{10}, K_{11}, L_{12}$ as positive and the numerator of $M_9, M_{10}, M_{11}, K_{12}$ as negative. After that, we obtain the following conditions,

$$\frac{2\lambda_1 \mu_2^2}{\mu_1^2} < \lambda_4 < \frac{2\lambda_2 \mu_1^2}{\mu_2^2} - \max\{\lambda_7, (\lambda_7 + 2\lambda_{10})\}, \quad (89)$$

$$\frac{2\lambda_3 \mu_2^2}{\mu_3^2} < \lambda_6 < \frac{2\lambda_2 \mu_3^2}{\mu_2^2} - \lambda_9, \quad (90)$$

$$\frac{2\lambda_1 \mu_3^2}{\mu_1^2} < (\lambda_5 + \lambda_8) < \frac{2\lambda_3 \mu_1^2}{\mu_3^2}, \quad (91)$$

where the inequalities (85) to (88) are derived from these new ones. And where the function max takes the largest value from a set.

Finally, we conclude under conditions above, the global minimum of the potential lies on point $\tilde{\mathbf{P}}_1$, where

$$\tilde{\mathbf{P}}_1 \cdot \tilde{g}_2 \cdot \tilde{\mathbf{P}}_1 = 4 K_{01} M_{01} > 0. \quad (92)$$

Also

$$\langle \underline{K} \rangle = \begin{pmatrix} \langle \Phi_1^\dagger \Phi_1 \rangle & \langle \Phi_3^\dagger \Phi_1 \rangle \\ \langle \Phi_1^\dagger \Phi_2 \rangle & \langle \Phi_2^\dagger \Phi_2 \rangle \end{pmatrix} = \begin{pmatrix} \frac{\lambda_5 \mu_3^2 - 2\lambda_3 \mu_2^2}{4\lambda_1 \lambda_3 - \lambda_5^2} & 0 \\ 0 & 0 \end{pmatrix}, \quad (93)$$

$$\langle \underline{L} \rangle = \begin{pmatrix} \langle \Phi_1^\dagger \Phi_1 \rangle & \langle \Phi_3^\dagger \Phi_1 \rangle \\ \langle \Phi_1^\dagger \Phi_3 \rangle & \langle \Phi_3^\dagger \Phi_3 \rangle \end{pmatrix} = \begin{pmatrix} \frac{\lambda_5 \mu_3^2 - 2\lambda_3 \mu_1^2}{4\lambda_1 \lambda_3 - \lambda_5^2} & 0 \\ 0 & \frac{\lambda_5 \mu_1^2 - 2\lambda_1 \mu_3^2}{4\lambda_1 \lambda_3 - \lambda_5^2} \end{pmatrix}, \quad (94)$$

$$\langle \underline{M} \rangle = \begin{pmatrix} \langle \Phi_2^\dagger \Phi_2 \rangle & \langle \Phi_3^\dagger \Phi_2 \rangle \\ \langle \Phi_2^\dagger \Phi_3 \rangle & \langle \Phi_3^\dagger \Phi_3 \rangle \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & \frac{\lambda_5 \mu_2^2 - 2\lambda_1 \mu_3^2}{4\lambda_1 \lambda_3 - \lambda_5^2} \end{pmatrix}, \quad (95)$$

which implies the configuration of VEVs vectors given by (27). Let us mention here that the former results does not change if we consider complex phases in the VEV. Then we have

$$\frac{v_1^2 + V_1^2}{2} = \frac{\lambda_5 \mu_3^2 - 2\lambda_3 \mu_1^2}{4\lambda_1 \lambda_3 - \lambda_5^2}, \quad (96)$$

$$\frac{v_3^2}{2} = \frac{\lambda_5 \mu_1^2 - 2\lambda_1 \mu_3^2}{4\lambda_1 \lambda_3 - \lambda_5^2}, \quad (97)$$

solutions that agree with the constraint equations given in (28). Using the former relations we can write the Lagrange multiplier u_1 in the following way

$$u_1 = \frac{1}{4} \frac{(4\lambda_1 \lambda_3 - \lambda_5^2)}{(\lambda_5 \mu_3^2 - 2\lambda_3 \mu_1^2)} \left(\frac{\lambda_4 (v_1^2 + V_1^2) + \lambda_6 v_3^2}{2} + \mu_2^2 \right) > 0. \quad (98)$$

Since the value in (96) is positive, the large parenthesis in (98) is also positive and thus, the square mass in (34) is

also positive. Using conditions (84), we can conclude that the square masses in (33), (36), (37) and (39) are also positive quantities, with the hierarchy $(M_{he3}^2, M_{ho1}^2, M_{h\pm}^2) > M_{he4}^2 = M_{ho2}^2$.

At the global minimum, the Higgs potential now becomes

$$V_{\min.} = \frac{1}{4} \mu_1^2 (v_1^2 + V_1^2) + \frac{1}{4} \mu_3^2 v_3^2 < 0. \quad (99)$$

Therefore, in order to have the deepest minimum value for the potential for this particular vacuum structure, the following conditions are highly suggested

$$v_1, V_1, v_3 \neq 0. \quad (100)$$

5.2 The general case $v_1, V_1, v_2, V_2 \neq 0$

In this case, we want the global minimum of potential be located at $\tilde{\mathbf{P}}_5$. Looking at the Table (2), we choose this point as the global minimum, because for it $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ are LD and the VEV configuration presents in Eqs. (4) and (5) are reproduced. For that, it is necessary that

$$\lambda_7 + 2\lambda_{10} < 0. \quad (101)$$

At the same time let us eliminate the possibility that the exceptional values v_7 and w_8 became global minima, which is reached by making them negatives, that is

$$\lambda_8, \lambda_9 > 0, \quad (102)$$

in such a way that the square mass in (25) becomes positive.

To exclude $\tilde{\mathbf{P}}_6$ as the global minimum it is sufficient to assume that $u_5 > u_6$ which is achieved as far as

$$\lambda_{10} < 0. \quad (103)$$

If we assume that

$$\lambda_4 < 0, \quad \lambda_5 < 0, \quad \text{and} \quad \lambda_6 < 0, \quad (104)$$

and taking into account (56), (57), (59), (65) and (101), it implies that $u_1 < 0, w_2 < 0, u_3 < 0, w_4 < 0, v_9 < 0, u_{13} < 0, v_{13} < 0, u_{14} < 0, w_{14} < 0, v_{15} < 0$ and $w_{15} < 0$. Finally, the remaining Lagrange multipliers $u_{10} < 0, v_{10} < 0, u_{11} < 0, v_{11} < 0, u_{12} < 0$ and $u_{12} < 0$ are negative, when the corresponding points, are stationary points respectively. From conditions (56), (59), (101), (102), (103) and (104), the inequalities (20), (23) and (25) are immediately satisfied.

As you can observe, being exhaustive in our reasoning, the global minimum remains at $\tilde{\mathbf{P}}_5$, and it is given by

$$\begin{pmatrix} K_5 = \frac{8 \mu_3^2 [w_{15} (u_5 - u_3) + v_{15} (u_5 - u_1)]}{d} \\ L_5 = \frac{8 \mu_3^2 [w_{15} (u_5 - u_3) + u_5^2] - 8 \mu_1^2 u_{13} w_4 - 2(2\lambda_4 \mu_3^2 - \lambda_5 \mu_2^2 - \lambda_6 \mu_1^2) u_5}{d} \\ M_5 = \frac{8 \mu_3^2 [v_{15} (u_5 - u_1) + u_5^2] - 8 \mu_1^2 u_{13} w_4 - 2(2\lambda_4 \mu_3^2 - \lambda_5 \mu_2^2 - \lambda_6 \mu_1^2) u_5}{d} \\ \frac{16 |\mu_3^2| \sqrt{(u_5 - u_1)(u_5 - u_3)} v_{15} w_{15}}{|d|} \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad (105)$$

with

$$d = \lambda_1 (4\lambda_2 \lambda_3 - \lambda_6^2) + \lambda_2 (4\lambda_1 \lambda_3 - \lambda_5^2) - \lambda_3 [4\lambda_1 \lambda_2 + (\lambda_4 - 4u_5)^2] + \lambda_5 \lambda_6 (\lambda_4 - 4u_5),$$

where the first two terms of d are positive and the last ones negative. In light of the results above, we can choose λ_1, λ_2 and λ_3 as larger as necessary such that

$$d > 0, \quad (106)$$

as can be observed from $\lim_{\lambda_{1,2,3} \rightarrow +\infty} d \approx 4\lambda_1 \lambda_2 \lambda_3 = \mathcal{O}(\lambda^3) > 0$ and similarly $\lim_{\lambda_{1,2,3} \rightarrow +\infty} L_5, M_5 = \mathcal{O}(1/\lambda) > 0$, such that it is possible to find cases for which

$$L_5 > 0, \quad \text{and} \quad M_5 > 0. \quad (107)$$

The fourth entry in (105) is taken positive by assuming positive VEV. Since we are in the domain given by (71b), we must have

$$\tilde{\mathbf{P}}_5 \cdot \tilde{g}_1 \cdot \tilde{\mathbf{P}}_5 = 0, \quad (108)$$

$$\tilde{\mathbf{P}}_5 \cdot \tilde{g}_2 \cdot \tilde{\mathbf{P}}_5 = -64 \mu_3^2 w_{15} (u_5 - u_3) [4 \mu_1^2 u_{13} w_4 - 4 \mu_3^2 u_5^2 - (\lambda_6 \mu_1^2 + \lambda_5 \mu_2^2 - 2\lambda_4 \mu_3^2) u_5] / d^2 > 0, \quad (109)$$

$$\tilde{\mathbf{P}}_5 \cdot \tilde{g}_3 \cdot \tilde{\mathbf{P}}_5 = -64 \mu_3^2 v_{15} (u_5 - u_1) [4 \mu_1^2 u_{13} w_4 - 4 \mu_3^2 u_5^2 - (\lambda_6 \mu_1^2 + \lambda_5 \mu_2^2 - 2\lambda_4 \mu_3^2) u_5] / d^2 > 0. \quad (110)$$

where we can observe too that $\lim_{\lambda_{1,2,3} \rightarrow +\infty} (\tilde{\mathbf{P}}_5 \cdot \tilde{g}_2 \cdot \tilde{\mathbf{P}}_5, \tilde{\mathbf{P}}_5 \cdot \tilde{g}_3 \cdot \tilde{\mathbf{P}}_5) = \mathcal{O}(1/\lambda^2) > 0$.

The expectation values satisfy also

$$\langle \underline{K} \rangle = \begin{pmatrix} \langle \Phi_1^\dagger \Phi_1 \rangle & \langle \Phi_1^\dagger \Phi_2 \rangle \\ \langle \Phi_1^\dagger \Phi_2 \rangle & \langle \Phi_2^\dagger \Phi_2 \rangle \end{pmatrix} \quad (111)$$

$$= \begin{pmatrix} \frac{8 \mu_3^2 w_{15} (u_5 - u_3)}{d} & \frac{8 |\mu_3^2| \sqrt{(u_5 - u_1)(u_5 - u_3) v_{15} w_{15}}}{|d|} \\ \frac{8 |\mu_3^2| \sqrt{(u_5 - u_1)(u_5 - u_3) v_{15} w_{15}}}{|d|} & \frac{8 \mu_3^2 v_{15} (u_5 - u_1)}{d} \end{pmatrix},$$

$$\langle \underline{L} \rangle = \begin{pmatrix} \langle \Phi_1^\dagger \Phi_1 \rangle & \langle \Phi_3^\dagger \Phi_1 \rangle \\ \langle \Phi_1^\dagger \Phi_3 \rangle & \langle \Phi_3^\dagger \Phi_3 \rangle \end{pmatrix} \quad (112)$$

$$= \begin{pmatrix} \frac{8 \mu_3^2 w_{15} (u_5 - u_3)}{d} & 0 \\ 0 & -\frac{2 [4 \mu_1^2 u_{13} w_4 - 4 \mu_3^2 u_5^2 - (\lambda_6 \mu_1^2 + \lambda_5 \mu_2^2 - 2\lambda_4 \mu_3^2) u_5]}{d} \end{pmatrix},$$

$$\langle \underline{M} \rangle = \begin{pmatrix} \langle \Phi_2^\dagger \Phi_2 \rangle & \langle \Phi_3^\dagger \Phi_2 \rangle \\ \langle \Phi_2^\dagger \Phi_3 \rangle & \langle \Phi_3^\dagger \Phi_3 \rangle \end{pmatrix} \quad (113)$$

$$= \begin{pmatrix} \frac{8 \mu_3^2 v_{15} (u_5 - u_1)}{d} & 0 \\ 0 & -\frac{2 [4 \mu_1^2 u_{13} w_4 - 4 \mu_3^2 u_5^2 - (\lambda_6 \mu_1^2 + \lambda_5 \mu_2^2 - 2\lambda_4 \mu_3^2) u_5]}{d} \end{pmatrix}.$$

This shows that $\langle \Phi_3 \rangle$ is orthogonal to $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$, and at the same time $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ are LD due to the relation (108). It also shows that generality is not lost by taking positive VEV in (4) and (5). We then have

$$\frac{v_1^2 + V_1^2}{2} = \frac{8 \mu_3^2 w_{15} (u_5 - u_3)}{d}, \quad (114)$$

$$\frac{v_2^2 + V_2^2}{2} = \frac{8 \mu_3^2 v_{15} (u_5 - u_1)}{d}, \quad (115)$$

$$\frac{v_3^2}{2} = -\frac{8 \mu_1^2 u_{13} w_4 - 8 \mu_3^2 u_5^2}{d} + \frac{2 (\lambda_6 \mu_1^2 + \lambda_5 \mu_2^2 - 2\lambda_4 \mu_3^2) u_5}{d}, \quad (116)$$

$$\frac{v_1 v_2 + V_1 V_2}{2} = \frac{8 |\mu_3^2| \sqrt{(u_5 - u_1)(u_5 - u_3) v_{15} w_{15}}}{|d|}, \quad (117)$$

The solution to (114), (115) and (116) coincide with the expressions given in (15). Also, from the relations (11), (114), (115) and (117) we can see that the VEV satisfy the following relations

$$\frac{v_1^2}{v_2^2} = \frac{V_1^2}{V_2^2} = \frac{w_{15} (u_5 - u_3)}{v_{15} (u_5 - u_1)} = \alpha^2, \quad (118)$$

where

$$\alpha = \sqrt{\frac{w_{15} (u_5 - u_3)}{v_{15} (u_5 - u_1)}} \quad (119)$$

is the proportionality factor between $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ as stated in (153), which allow us to connect the value of α to the Lagrange multipliers.

At the global minimum the Higgs potential becomes

$$V_{\min.} = \frac{1}{4} \mu_1^2 (v_1^2 + V_1^2) + \frac{1}{4} \mu_2^2 (v_2^2 + V_2^2) + \frac{1}{4} \mu_3^2 v_3^2 < 0, \quad (120)$$

which reproduces Eq. (99) in the limit $v_2 = V_2 = 0$. Therefore, in order to have the deepest minimum value for the potential for this particular vacuum structure, the following conditions are highly suggested

$$v_1, V_1, v_2, V_2, v_3 \neq 0. \quad (121)$$

6 Conclusions

In this paper we have presented original results related to the scalar sector of some 3-3-1 models without exotic electric charges. An exhaustive study of the scalar potential with 3 scalar triplets Φ_1, Φ_2 and Φ_3 and VEV as introduced in Sects. (2.3.1) and (2.3.2); potential which does not include the possible cubic term according to the discrete symmetry $\Phi_1 \rightarrow -\Phi_1$ imposed, has been carried through. This problem partially analyzed in the literature [7, 14, 15] had not been studied in a systematic way.

In concrete we have:

- Looked for a consistent implementation of the Higgs mechanism.
- Implemented a consistent electroweak symmetry breaking pattern.
- Established the strong stability conditions for the scalar potential.
- Found the stationary points of the scalar potential, except the ones coming from specific relations among the parameters μ_1^2, μ_2^2 and μ_3^2 that we assume are not satisfied, in general.

One outstanding new result is that $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$, the VEV of the two Higgs scalars with identical quantum numbers, must be proportional to each other, a necessary condition in order to properly implement the Higgs mechanism, and achieve a consistent electroweak symmetry breaking; besides, the proportionality constant is connected with the Lagrange multipliers (which in turn are connected to the other parameters of the potential) via Eq. (119).

Other important result is that, from the nine possible vacuum structures compatible with the stated LD constraint, only two are independent. Our analysis has been done for both structures.

But probably, the most important conclusion of our study is the existence (as a sufficient condition), in the scalar potential, of a global minimum stationary point for each one of the two cases of VEV considered, being it, at the same time, compatible with the stability conditions imposed in the strong sense. Stability conditions and the global minimum point were found via the orbit gauge method, implemented in this case for three scalar triplets. Although the method can give unphysical VEVs.

Acknowledgments

We both acknowledge the warm hospitality from the “Laboratorio de Física Teórica de la Universidad de La Plata, in La Plata, Argentina”, during several stages of the development of the present work. Also, very illuminating were several conversations with professors C. García-Canal, D. Gómez-Dumm and H. Fanchiotti. We thank in special C. García-Canal for a critical reading of the original manuscript. Y.G. acknowledges financial support and leave of absence from “Universidad de Nariño” in Pasto, Colombia. The present work is part of the doctoral thesis of Y.G.

A A review of the algebraic method used

Let us briefly review in this appendix, and following Refs. [3] and [4], a new algebraic approach used to determine the global minimum of the Higgs scalar potential, its stability, and the spontaneous symmetry breaking from $SU(2)_L \otimes U(1)_Y$ down to $U(1)_{EM}$, in the extension of the SM known as the THDM. This mathematical approach and its notation was used in Secs (4) and (5) of this paper, generalizing it to the study of the scalar sector of some 3-3-1 models.

Stability, and the stationary points of the scalar potential for the THDM can be analyzed in terms of four real quantities given by

$$K_0 = \sum_{i=1,2} \varphi_i^\dagger \varphi_i, \quad K_a = \sum_{i,j=1,2} (\varphi_i^\dagger \varphi_j) \sigma_{ij}^a, \quad (a = 1, 2, 3). \quad (122)$$

where φ_1 and φ_2 stand for two Higgs scalar field doublets with identical quantum numbers and $\sigma^a (a = 1, 2, 3)$ are the Pauli spin matrices. The four vector (K_0, \mathbf{K}) must lie on or inside the forward *light cone*, that is

$$K_0 \geq 0, \quad K_0^2 - \mathbf{K}^2 \geq 0. \quad (123)$$

Then the positive and hermitian 2×2 matrix

$$\underline{K} = \begin{pmatrix} \varphi_1^\dagger \varphi_1 & \varphi_2^\dagger \varphi_1 \\ \varphi_1^\dagger \varphi_2 & \varphi_2^\dagger \varphi_2 \end{pmatrix} \quad (124)$$

may be written as

$$\underline{K}_{ij} = \frac{1}{2} (K_0 \delta_{ij} + K_a \sigma_{ij}^a). \quad (125)$$

Inverting Eq. (122) one obtains

$$\begin{aligned} \varphi_1^\dagger \varphi_1 &= (K_0 + K_3)/2, & \varphi_1^\dagger \varphi_2 &= (K_1 + iK_2)/2, \\ \varphi_2^\dagger \varphi_2 &= (K_0 - K_3)/2, & \varphi_2^\dagger \varphi_1 &= (K_1 - iK_2)/2. \end{aligned} \quad (126)$$

The most general $SU(2)_L \otimes U(1)_Y$ invariant Higgs scalar potential can thus be expressed as

$$V(\varphi_1, \varphi_2) = V_2 + V_4; \quad (127a)$$

$$V_2 = \xi_0 K_0 + \xi_a K_a, \quad (127b)$$

$$V_4 = \eta_{00} K_0^2 + 2K_0 \eta_a K_a + K_a \eta_{ab} K_b, \quad (127c)$$

where the 14 independent parameters $\xi_0, \xi_a, \eta_{00}, \eta_a$ and $\eta_{ab} = \eta_{ba}$ are real. Subsequently, it is defined $\mathbf{K} = (K_a)$, $\boldsymbol{\xi} = (\xi_a)$, $\boldsymbol{\eta} = (\eta_a)$ and $E = (\eta_{ab})$.

A.1 Stability

From (127), for $K_0 > 0$ and defining $\mathbf{k} = \mathbf{K}/K_0$, one obtains

$$V_2 = K_0 J_2(\mathbf{k}), \quad J_2(\mathbf{k}) := \xi_0 + \boldsymbol{\xi}^T \mathbf{k}, \quad (128)$$

$$V_4 = K_0^2 J_4(\mathbf{k}), \quad J_4(\mathbf{k}) := \eta_{00} + 2\boldsymbol{\eta}^T \mathbf{k} + \mathbf{k}^T E \mathbf{k}, \quad (129)$$

where the functions $J_2(\mathbf{k})$ and $J_4(\mathbf{k})$ on the domain $|\mathbf{k}| \leq 1$ have been introduced.

For the potential to be stable, it must be bounded from below. The stability is determined by the behavior of $V = V_1 + V_2$ in the limit $K_0 \rightarrow \infty$, hence by the signs of $J_2(\mathbf{k})$ and/or $J_4(\mathbf{k})$ in (128) and (129). In the analysis, only the *strong* criterion for stability is analyzed. That is, the stability is going to be determined solely by V_4 (the V quartic term). That is, we demand that

$$J_4(\mathbf{k}) > 0 \quad \text{for all } |\mathbf{k}| \leq 1. \quad (130)$$

To assure that $J_4(\mathbf{k})$ is always positive, it is sufficient to consider its value for all its stationary points on the domain $|\mathbf{k}| < 1$ and for all the stationary points on the boundary $|\mathbf{k}| = 1$. This leads to bounds on the parameters η_{00}, η_a and η_{ab} of the quartic term V_4 of the scalar potential. The regular solutions are obtained by inverting the matrix $(E - u)$ which appears in the calculations, where u is the Lagrange multiplier associated to the boundary condition under consideration. When the matrix is not invertible, we say that we have an exceptional solution. The two cases $|\mathbf{k}| < 1$ and $|\mathbf{k}| = 1$ lead to the functions

$$f(u) = u + \eta_{00} - \boldsymbol{\eta}^T (E - u)^{-1} \boldsymbol{\eta}, \quad (131)$$

$$f'(u) = 1 - \boldsymbol{\eta}^T (E - u)^{-2} \boldsymbol{\eta}, \quad (132)$$

so that for all “regular” stationary points \mathbf{k} of $J_4(\mathbf{k})$

$$f(u) = J_4(\mathbf{k})|_{stat}, \quad \text{and} \quad (133)$$

$$f'(u) = 1 - \mathbf{k}^2 \quad (134)$$

hold, where $u = 0$ must be set for the solution with $|\mathbf{k}| < 1$. There are stationary points of $J_4(\mathbf{k})$ with $|\mathbf{k}| < 1$ and

$|\mathbf{k}| = 1$ exactly if $f'(0) > 0$ and $f'(u) = 0$, respectively, and the value of $J_4(\mathbf{k})$ is then given by $f(u)$.

In a basis where $E = \text{diag}(\mu_1, \mu_2, \mu_3)$ it is obtained

$$f(u) = u + \eta_{00} - \sum_{a=1}^3 \frac{\eta_a^2}{\mu_a - u}, \quad (135)$$

$$f'(u) = 1 - \sum_{a=1}^3 \frac{\eta_a^2}{(\mu_a - u)^2}. \quad (136)$$

The derivative $f'(u)$ has at most 6 zeros. Notice that there are no exceptional solutions if in this basis all three components of $\boldsymbol{\eta}$ are different from zero.

Consider now the functions $f(u)$ and $f'(u)$ and denote by I

$$I = \{u_1, \dots, u_n\}, \quad (137)$$

the set of values u_j for which $f'(u_j) = 0$. Add $u_k = 0$ to I if $f'(0) > 0$. Consider then the eigenvalues μ_a ($a = 1, 2, 3$) of E . Add those μ_a to I where $f(\mu_a)$ is finite and $f'(\mu_a) \geq 0$. Then $n \leq 10$. The values of the function $J_4(\mathbf{k})$ at its stationary points are given by

$$J_4(\mathbf{k})|_{\text{stat}} = f(u_i) \quad (138)$$

with $u_i \in I$. In Ref. [13] it was shown that the stationary point in I having the smallest value, will produce the smallest value of $J_4(\mathbf{k})$ in the domain $|\mathbf{k}| \leq 1$. Then, one states the following theorem

Theorem 1 *The global minimum of the function $J_4(\mathbf{k})$, in the domain $|\mathbf{k}| \leq 1$, is given and guaranteed by the stationary point of the set I with the smallest value.*

Namely, this result guarantees strong stability if $f(u) > 0$, where u is the smallest value of I . The potential is unstable if we have $f(u) < 0$. If $f(u) = 0$ we have to consider in addition $J_2(\mathbf{k})$ in order to decide on the stability of the potential.

A.2 Stationary points

After the stability analysis is done, the next step is to determine the location of the stationary points of the scalar potential, since among these points the local and global minima are found. To this end it is defined

$$\tilde{\mathbf{K}} = \begin{pmatrix} K_0 \\ \mathbf{0} \end{pmatrix}, \quad \tilde{\boldsymbol{\xi}} = \begin{pmatrix} \xi_0 \\ \boldsymbol{\xi} \end{pmatrix}, \quad \tilde{E} = \begin{pmatrix} \eta_{00} & \boldsymbol{\eta}_E^T \\ \boldsymbol{\eta} & E \end{pmatrix}. \quad (139)$$

In this notation the potential (127) reads

$$V = \tilde{\mathbf{K}}^T \tilde{\boldsymbol{\xi}} + \tilde{\mathbf{K}}^T \tilde{E} \tilde{\mathbf{K}} \quad (140)$$

and is defined on the domain

$$\tilde{\mathbf{K}}^T \tilde{g} \tilde{\mathbf{K}} \geq 0, \quad K_0 \geq 0, \quad (141)$$

with

$$\tilde{g} = \begin{pmatrix} 1 & \mathbf{0} \\ \mathbf{0} & -\mathbb{1} \end{pmatrix}. \quad (142)$$

For the discussion of the stationary points of V , three different cases must be distinguished: $\tilde{\mathbf{K}} = 0$, $K_0 > |\mathbf{K}|$ which are the solutions inside the forward light cone, and $K_0 = |\mathbf{K}| > 0$ which are the solutions on the forward light cone.

The trivial configuration $\tilde{\mathbf{K}} = 0$ is a stationary point of the potential with $V = 0$, as a direct consequence of the definitions. The stationary points of V in the inner part of the domain, $K_0 > |\mathbf{K}|$, are given by

$$\tilde{E} \tilde{\mathbf{K}} = -\frac{1}{2} \tilde{\boldsymbol{\xi}}, \quad \text{with } \tilde{\mathbf{K}}^T \tilde{g} \tilde{\mathbf{K}} > 0 \quad \text{and } K_0 > 0. \quad (143)$$

The stationary points of V on the domain boundary $K_0 = |\mathbf{K}| > 0$ are stationary points of the function

$$\tilde{F}(\tilde{\mathbf{K}}, w) = V - w \tilde{\mathbf{K}}^T \tilde{g} \tilde{\mathbf{K}}, \quad (144)$$

where w is a Lagrange multiplier. The relevant stationary points of \tilde{F} are given by

$$(\tilde{E} - w \tilde{g}) \tilde{\mathbf{K}} = -\frac{1}{2} \tilde{\boldsymbol{\xi}}, \quad \text{with } \tilde{\mathbf{K}}^T \tilde{g} \tilde{\mathbf{K}} = 0 \quad \text{and } K_0 > 0. \quad (145)$$

For *any* stationary point, the potential is given by

$$V|_{\text{stat}} = \frac{1}{2} \tilde{\mathbf{K}}^T \tilde{\boldsymbol{\xi}} = -\tilde{\mathbf{K}}^T \tilde{E} \tilde{\mathbf{K}}. \quad (146)$$

Similarly to the stability analysis in Sec. A.1, a unified description for the regular stationary points of V with $K_0 > 0$ for both $|\mathbf{K}| < K_0$ and $|\mathbf{K}| = K_0$ can be used by defining the functions

$$\tilde{f}(w) = -\frac{1}{4} \tilde{\boldsymbol{\xi}}^T (\tilde{E} - w \tilde{g})^{-1} \tilde{\boldsymbol{\xi}}, \quad (147)$$

$$\tilde{f}'(w) = -\frac{1}{4} \tilde{\boldsymbol{\xi}}^T (\tilde{E} - w \tilde{g})^{-1} \tilde{g} (\tilde{E} - w \tilde{g})^{-1} \tilde{\boldsymbol{\xi}}. \quad (148)$$

Denoting the first component of $\tilde{\mathbf{K}}(w)$ as $K_0(w)$ the following theorem holds

Theorem 2 *The stationary points of the potential are given by*

- (Ia) $\tilde{\mathbf{K}} = \tilde{\mathbf{K}}(0)$ if $\tilde{f}'(0) < 0$, $K_0(0) > 0$ and $\det \tilde{E} \neq 0$,
- (Ib) solutions $\tilde{\mathbf{K}}$ of (143) if $\det \tilde{E} = 0$,
- (IIa) $\tilde{\mathbf{K}} = \tilde{\mathbf{K}}(w)$ for w with $\det(\tilde{E} - w \tilde{g}) \neq 0$, $\tilde{f}'(w) = 0$ and $K_0(w) > 0$,
- (IIb) solutions $\tilde{\mathbf{K}}$ of (145) for w with $\det(\tilde{E} - w \tilde{g}) = 0$,
- (III) $\tilde{\mathbf{K}} = 0$.

In what follows it is assumed that the potential is stable. For parameters fulfilling $\xi_0 \geq |\boldsymbol{\xi}|$, this immediately implies $J_2(\mathbf{k}) \geq 0$ and hence, from the strong condition (130), $V > 0$ for all $\tilde{\mathbf{K}} \neq 0$. Therefore for these parameters the global minimum is at $\tilde{\mathbf{K}} = 0$. This leads to the requirement

$$\xi_0 < |\boldsymbol{\xi}|. \quad (149)$$

Also, it is obtained

$$\left. \frac{\partial V}{\partial K_0} \right|_{\substack{\mathbf{k} \text{ fixed,} \\ K_0=0}} = \xi_0 + \boldsymbol{\xi}^T \mathbf{k} < 0 \quad (150)$$

for some \mathbf{k} , i.e. the global minimum of V lies at $\tilde{\mathbf{K}} \neq 0$ with

$$V|_{min} < 0. \quad (151)$$

Firstly, consider $p_0 = |\mathbf{p}|$. From (140) and (145) it follows

$$\left. \frac{\partial V}{\partial K_0} \right|_{\substack{\mathbf{K} \text{ fixed,} \\ \tilde{\mathbf{K}} = \tilde{\mathbf{p}}}} = \xi_0 + 2(\tilde{E}\tilde{\mathbf{p}})_0 = 2w_p p_0. \quad (152)$$

If $w_p < 0$, there are points $\tilde{\mathbf{K}}$ with $K_0 > p_0$, $\mathbf{K} = \mathbf{p}$ and lower potential in the neighborhood of $\tilde{\mathbf{p}}$, which therefore cannot be a minimum. The conclusion is that in a theory with the required electroweak symmetry breaking (EWSB) the global minimum must have a Lagrange multiplier such that $w_0 \geq 0$, and for the THDM, the global minimum lies on the stationary points of the classes (IIa) and (IIb) of theorem 2, with the largest Lagrange multiplier [3].

B Linear dependence between $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$

In this appendix we study the consequences of a linear dependence between $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$.

As in the main text we use the VEV

$$\langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \\ V_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 \\ V_2 \end{pmatrix},$$

$$\langle \Phi_3 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v_3 \\ 0 \\ 0 \end{pmatrix}.$$

The LD between $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ can be written as

$$\langle \Phi_1 \rangle = \alpha \langle \Phi_2 \rangle, \quad (153)$$

where α is a constant. Eq. (153) implies that $v_1 = \alpha v_2$ and $V_1 = \alpha V_2$, which combine to produce the constraint $v_2 V_1 = v_1 V_2$.

Now, the nine $U(3)$ generators are

$$\begin{aligned} I_3 &= \sqrt{\frac{2}{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \lambda_3 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad (154) \\ \lambda_6 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \end{aligned}$$

where $\lambda_i (i = 1, \dots, 8)$ are the eight Gell-Mann unitary matrices for $SU(3)$.

Let us now show that the LD in Eq. (153) with the additional constraint $\langle \Phi_3 \rangle \neq 0$, implies that either I_3 , or a linear combination of the generators in (154) which includes I_3 , remains unbroken, with the consequence that the appearance of an extra zero mass Goldstone Bosons is avoided.

The algebra shows that the most general new unbroken generator is given by the following linear combination:

$$G = aI_3 + b\lambda_3 + c\lambda_8 + d\lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & V_1^2 & -v_1 V_1 \\ 0 & -v_1 V_1 & v_1^2 \end{pmatrix}, \quad (155)$$

where

$$\begin{aligned} a &= \frac{V_1^2 + v_1^2}{2}, & b &= -\frac{V_1^2}{2}, \\ c &= \frac{V_1^2 - 2v_1^2}{2\sqrt{3}}, & d &= -v_1 V_1. \end{aligned}$$

That G remains unbroken can be seen by the fact that $G\langle \Phi_1 \rangle = 0$ by direct calculation, $G\langle \Phi_2 \rangle = 0$ is a consequence of the relation (11), and $G\langle \Phi_3 \rangle = 0$ is trivial.

Since $Tr.G = v_1^2 + V_1^2 \neq 0$, the new unbroken generator in Eq. (155) is such that $G \in U(3)$ but $G \notin SU(3)$.

C Discrete symmetry in the scalar potential

Under assumption of the discrete symmetry $\Phi_1 \rightarrow -\Phi_1$, the most general potential obtained from (9), can then be written in the following form:

$$\begin{aligned} V(\Phi_1, \Phi_2, \Phi_3) &= \mu_1^2 \Phi_1^\dagger \Phi_1 + \mu_2^2 \Phi_2^\dagger \Phi_2 + \mu_3^2 \Phi_3^\dagger \Phi_3 + \lambda_1 (\Phi_1^\dagger \Phi_1)^2 \\ &+ \lambda_2 (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_3^\dagger \Phi_3)^2 + \lambda_4 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\ &+ \lambda_5 (\Phi_1^\dagger \Phi_1) (\Phi_3^\dagger \Phi_3) + \lambda_6 (\Phi_2^\dagger \Phi_2) (\Phi_3^\dagger \Phi_3) + \lambda_7 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ &+ \lambda_8 (\Phi_1^\dagger \Phi_3) (\Phi_3^\dagger \Phi_1) + \lambda_9 (\Phi_2^\dagger \Phi_3) (\Phi_3^\dagger \Phi_2) + (f \Phi_1^\dagger \Phi_2 + f^* \Phi_2^\dagger \Phi_1)^2. \end{aligned} \quad (156)$$

where the complex value f is going to be used as $f = f_1 + if_2$, with f_j , $j = 1, 2$ are two real parameters. With the new definitions of the scalar fields introduced in (12), and by demanding that the VEV in (4) and (5) became stationary points of the potential, the following nine constraints must be satisfied

$$\begin{aligned} \left. \frac{\partial V}{\partial H_1} \right|_{\text{fields}=0} &= \\ &[2\mu_1^2 v_1 + \lambda_4 v_1 V_2^2 + (\lambda_7 + 4f_1^2) v_2 V_1 V_2 + 2\lambda_1 v_1 V_1^2 \\ &+ (\lambda_7 + \lambda_4 + 4f_1^2) v_1 v_2^2 + 2\lambda_1 v_1^3 + \lambda_5 v_3^2 v_1] / 2 = 0, \end{aligned} \quad (157)$$

$$\begin{aligned} \left. \frac{\partial V}{\partial H_1'} \right|_{\text{fields}=0} &= \\ &[2\mu_1^2 V_1 + (\lambda_7 + \lambda_4 + 4f_1^2) V_1 V_2^2 + (\lambda_7 + 4f_1^2) v_1 v_2 V_2 \\ &+ 2\lambda_1 V_1^3 + (\lambda_4 v_2^2 + 2\lambda_1 v_1^2 + \lambda_5 v_3^2) V_1] / 2 = 0, \end{aligned} \quad (158)$$

$$\begin{aligned} \left. \frac{\partial V}{\partial H_2} \right|_{\text{fields}=0} &= \\ &\{2\mu_2^2 v_2 + 2\lambda_2 v_2 V_2^2 + (\lambda_7 + 4f_1^2) v_1 V_1 V_2 + \lambda_4 v_2 V_1^2 \\ &+ 2\lambda_2 v_2^3 + [(\lambda_7 + \lambda_4 + 4f_1^2) v_1^2 + \lambda_6 v_3^2] v_2\} / 2 = 0, \end{aligned} \quad (159)$$

$$\begin{aligned} \left. \frac{\partial V}{\partial H_2'} \right|_{\text{fields}=0} &= \\ &\{2\mu_2^2 V_2 + 2\lambda_2 V_2^3 + (\lambda_7 + 4f_1^2) v_1 v_2 V_1 \\ &+ [(\lambda_7 + \lambda_4 + 4f_1^2) V_1^2 + 2\lambda_2 v_2^2 + \lambda_4 v_1^2 + \lambda_6 v_3^2] V_2\} / 2 = 0, \end{aligned} \quad (160)$$

$$\left. \frac{\partial V}{\partial H_3} \right|_{\text{fields}=0} = \frac{v_3 (2\mu_3^2 + \lambda_6 V_2^2 + \lambda_5 V_1^2 + \lambda_6 v_2^2 + \lambda_5 v_1^2 + 2\lambda_3 v_3^2)}{2} = 0, \quad (161)$$

$$\left. \frac{\partial V}{\partial A_1} \right|_{\text{fields}=0} = 2f_1 f_2 v_2 (V_1 V_2 + v_1 v_2) = 0, \quad (162)$$

$$\left. \frac{\partial V}{\partial A'_1} \right|_{\text{fields}=0} = 2f_1 f_2 V_2 (V_1 V_2 + v_1 v_2) = 0, \quad (163)$$

$$\left. \frac{\partial V}{\partial A_2} \right|_{\text{fields}=0} = -2f_1 f_2 v_1 (V_1 V_2 + v_1 v_2) = 0, \quad (164)$$

$$\left. \frac{\partial V}{\partial A'_2} \right|_{\text{fields}=0} = -2f_1 f_2 V_1 (V_1 V_2 + v_1 v_2) = 0. \quad (165)$$

A simple algebra shows that both operations [$v_1 \times (158) - V_1 \times (157)$] and [$V_2 \times (159) - v_2 \times (160)$] produce the same relation

$$(\lambda_7 + 4f_1^2)(v_1 V_2 - v_2 V_1)(V_1 V_2 + v_1 v_2) = 0, \quad (166)$$

which must be satisfied in order to have a consistent set of equations (157)-(160).

The two possible solutions to (166) are $(v_1 V_2 - v_2 V_1) = 0$ and/or $(V_1 V_2 + v_1 v_2) = 0$. Obviously, (166) is satisfied if $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ are LD. (For the unphysical case $\langle \Phi_3 \rangle = 0$ with $\langle \Phi_1 \rangle$ and $\langle \Phi_2 \rangle$ being linearly independent, the mathematical solution $V_1 V_2 = -v_1 v_2$ is still available.)

But at the same time, the relations (162)-(165) must be satisfied, the alternative which remains for the physical case is that either the real or the imaginary part of f become zero, that is

$$f_1 = 0 \quad \text{or} \quad f_2 = 0, \quad (167)$$

meaning that f represents only one parameter, something which allow us to introduce the usual notation $|f|^2 = \frac{\lambda_{10}}{2}$, with λ_{10} either positive or negative.

References

1. Author, Journal **Volume**, (year) page numbers.
2. Author, *Book title* (Publisher, place year) page numbers
1. N.G. Deshpande and E. Ma, Phys. Rev. D **18**, 2574 (1978); H.E.haber, G.L.Kane and T.Sterling, Nucl. Phys. **B161**, 493 (1979); J.F.Donoghue and L.F.Li, Phys. Rev. **D19**, 945 (1979); L.J.Hall and M.B.Wise, Nucl. Phys. **B187**, 397 (1981); T.P.Cheng and M.Sher, Phys. Rev. **D35**, 3484 (1987).
2. J.F. Gunion, H.E. Haber, G. Kane, S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Reading, 1990); R.A. Diaz Sánchez, Ph. D. Thesis, hep-ph/0212237; I.F. Ginzburg and M. Krawczyk, Phys. Rev. D **72** (2005) 115013; S. Davidson and H.E. Haber, Phys. Rev. D **72** (2005) 035004 [Erratum-ibid. D **72** (2005) 099902]; H.E. Haber and D. O'Neil, Phys. Rev. D **74** (2006) 015018 [Erratum-ibid. D **74** (2006) 059905]; A. Barroso, P.M. Ferreira and R. Santos, Phys. Lett. B **632**, 684 (2006).
3. M. Maniatis, M.A. Manteuffel, O. Nachtmann and F. Nagel, Eur. Phys. J. C **48**, 805 (2006); M. Maniatis, M.A. Manteuffel and O. Nachtmann, Eur. Phys. J. C **48**, 1067 (2007).
4. F. Nagel, *New aspects of gauge-boson couplings and the Higgs sector*, PhD-thesis, Heidelberg University (2004).
5. I.P.Ivanov, Phys. Lett. **B632**, 360 (2006); I.P.Ivanov, Phys. Rev. **D75**, 035001 (2007) [Erratum-ibid **D76**. 039902 (2007)]; C.C.Nishi, Phys. Rev. **D74**. 036003 (2006); C.C.Nishi, Phys. Rev. **D77**, 055009 (2008).
6. F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); P.H. Frampton, Phys. Rev. Lett. **69**, 2887 (1992); V. Pleitez and M.D. Tonasse, Phys. Rev. D **48**, 2353 (1993); *ibid* 5274 (1993); D. Ng, Phys. Rev. D **49**, 4805 (1994); L. Epele, H. Fanchiotti, C. García Canal and D. Gómez Dumm, Phys. Lett. B **343** 291 (1995); M. Özer, Phys. Rev. D **54**, 4561 (1996).
7. J.C. Montero, F.Pisano and V. Pleitez, Phys. Rev. D **47**, 2918 (1993); R. Foot, H.N. Long and T.A. Tran, Phys. Rev. D **50**, R34 (1994); H.N. Long, Phys. Rev. D **53**, 437 (1996); *ibid* **54**, 4691 (1996).
8. V. Pleitez, Phys. Rev. D **53**, 514 (1996); M. Özer, Phys. Rev. D **54**, 1143, (1996); W.A.Ponce and O.Zapata, Phys. Rev. **D74**, 093007 (2006); J.C.Salazar, W.A.Ponce and D.A.Gutiérrez, Phys. Rev. **D75**, 075016 (2007).
9. D.L.Anderson and M.Sher, Phys. Rev. **D72**, 095014 (2005).
10. W.A.Ponce, Y.Giraldo, and L.A.Sánchez, *Systematic Study of 3-3-1 models*, in Proceedings of the VIII Mexican Workshop of Particles and Fields, Zacatecas, México, 2001, edited by J.L.Díaz-Cruz *et al*, AIP Conf. Proc. No. 623 (AIP, New York, 2002), pp. 341-346; W.A.Ponce, J.B.Flórez, and L.A.Sánchez, Int. J. Mod. Phys.**A17**, 643 (2002).
11. M.D.Tonasse, Phys. Lett. **B381**, 191 (1996); D.Gómez Dumm, Int. J. Mod. Phys. **A13**, 887 (1996); H.N.Long, Mod. Phys. Lett. **A13**, 1865 (1998); N.T.Anh, N.A.Ky and H.N.Long, Int. J. Mod. Phys. **A15**, 283 (2000); **16**, 541 (2001); M.B.Tully, G.C.Joshi, Int. J. Mod. Phys. **A18**, 1573 (2003).
12. W.A. Ponce, Y. Giraldo and L.A. Sánchez, Phys. Rev. D **67**, 075001 (2003).
13. Y. Giraldo, W.A.Ponce and L.A.Sánchez, Eur. Phys. J. **C63**, 461 (2009).
14. H.N.Long, Mod. Phys. Lett. **A13**, 1865 (1998); H.N.Long and N.Q.Lan, Europhys. Lett **64**, 571 (2003); J.L. García Luna, G. Tavares-Velasco and J.J. Toscano, Phys. Rev. **D69**. 093005 (2004); Alex G. Dias, C.A. de S.Pires and P.S.Rodrigues da Silva, Phys. Lett. **B628**, 85 (2005).
15. Palash B. Pal, Phys. Rev.**D52**,1659 (1995); Alex G. Dias, C.A. de S. Pires and P.S. Rodrigues da Silva, Phys.Rev.**D68**,115009 (2003); C.A. de S.Pires and P.S. Rodrigues da Silva, Eur.Phys.J.**C36**,397 (2004); Alex G. Dias, C.A. de S.Pires, V.Pleitez and P.S.Rodrigues da Silva, Phys. Lett. **B621**, 151 (2005).
16. P.V. Dong, H.N. Long, D.T. Nhung, and D.V. Soa, Phys. Rev. D **73**, 035004 (2006); P.V. Dong, H.N. Long, and D.V. Soa, Phys. Rev. D **73**, 075005 (2006); P.V.Dong, H.N.Long and D.T. Nhung, Phys. Lett **B639**, 527 (2006); D.A.Gutierrez, W.A.Ponce and L.A.Sánchez, Int. J. Mod. Phys. **A21**, 2217 (2006); P.V.Dong, H.N.Long and D.V.Soa, Phys. Rev. **D75**, 073006 (2007); P.V. Dong and H.N. Long, Adv. High Energy Phys. **2008**, 739492 (2008); J.M.Cabarcas, D.Gómez-Dumm and R.Martinez, Phys. Rev. **D77**, 036002 (2008).
17. P.V.Dong, D.T.Huong, Tr.T.Huong and H.N.Long, Phys. Rev. **D74**, 053003 (2006).
18. H. Fanchiotti, C. García-Canal, and W.A. Ponce, Eur. Phys. Lett. **72**, 733 (2005); J.C. Salazar, W.A. Ponce and D.A. Gutiérrez, Phys. Rev. **D75**, 075016 (2007).

19. R.Díaz, R.Martínez and F.Ochoa, Phys. Rev. **D69**, 095009 (2004); S.Filippi, W.A.Ponce and L.A.Sánchez, Eur. Phys. Lett. **73**, 142 (2006).
20. R.D.Peccei and H.R.Quinn, Phys. Rev. Lett. **38**, 1440 (1977); Phys. Rev. **D16**, 1791 (1977).