



Departamento de Física
Universidad de Nariño
Calle 18 No. 50-02
San Juan de Pasto – Colombia
Phone: +57-2-7244309

Grupo de Física de altas Energías
GFAE

[http://facultades.udenar.edu.co/facultad-de-ciencias-exactas-y-naturales/nuestros-docentes/docentes-fisica/rojas@udenar.edu.co](http://facultades.udenar.edu.co/facultad-de-ciencias-exactas-y-naturales/nuestros-docentes/docentes-fisica-rojas@udenar.edu.co)
eduro4000@gmail.com

San Juan de Pasto, 13 de Junio de 2020

Señores:

Asamblea de profesores

Departamento de física
Universidad de Nariño

Por medio de la presente informo de las actividades desarrolladas durante mi comisión académica en el evento **Workshop Materia Oscura en Colombia (MOCa 2022)**, que tuvo lugar en Bogotá del 31 de mayo al 1 de junio del 2022.

- Presenté la charla: ***A dark matter connection in a flavored axion model.***
- Y participé de las actividades realizadas en esta conferencia, que incluyeron: charlas principales y sesiones paralelas.
- Para la realización de este evento fui parte del comité organizador como representante de la universidad de Nariño.

El resumen de mi charla es: A Peccei-Quinn (PQ) symmetry is proposed, in order to generate in the Standard Model (SM) quark sector a realistic mass matrix ansatz with five texture-zeros. Limiting our analysis to Hermitian mass matrices we show that this requires a minimum of 4 Higgs doublets. This model allows assigning values close to 1 for several Yukawa couplings, giving insight into the origin of the mass scales in the SM. Since the PQ charges are non-universal the model features Flavor-Changing Neutral Currents (FCNC) at the tree level. In the model We found a pseudoscalar Nambu-Goldstone which could be a dark matter candidate.

EDUARDO ROJAS PEÑA Ph.D

Profesor Tiempo Completo
Departamento de física
Universidad de Nariño

A dark matter connection in a flavored axion model

Presented by Eduardo Rojas

In collaboration with: Y. Giraldo, R. Martínez, J.C. Salazar

Universidad de Nariño



eduro4000@gmail.com

June 1, 2022

Outline

- 1 Strong CP problem
- 2 The PQ solution.
- 3 The hierarchy problem
- 4 The model particle content.
- 5 Low energy constraints
- 6 A dark matter candidate
- 7 Conclusions

The $U(1)_A$ problem

- In the limit where the SM fermions are massless the QCD lagrangian has the symmetry $U(N)_V \otimes U(N)_A$, for the first family $N = 2$.
 $U(2)_V = SU(2)_V \otimes U(1)_V = SU(2)_I$ isospin symmetry $\otimes U(1)_B$ Baryon number conservation.

The $U(1)_A$ problem

- In the limit where the SM fermions are massless the QCD lagrangian has the symmetry $U(N)_V \otimes U(N)_A$, for the first family $N = 2$.
 $U(2)_V = SU(2)_V \otimes U(1)_V = SU(2)_I$ isospin symmetry $\otimes U(1)_B$ Baryon number conservation.
- The global $U(2)_A = SU(2)_A \otimes U(1)_A$ symmetry is broken spontaneously by the quark condensate, thus, one expects 4 Nambu-Goldstone bosons ($\pi^0, \pi^-, \pi^+, \eta(?)$), but η is too heavy. Although pions are light, there is no clue of another light state in the hadronic spectrum. Weinberg dubbed this the $U(1)_A$ problem, suggesting that, somehow, there was no $U(1)_A$ symmetry in QCD.

The Strong CP problem

1. 'T Hooft realized that the current associated with the $U(1)_A$ symmetry is anomalous, i.e., $\partial_\mu J_5^\mu = \frac{g^2 N}{32\pi^2} F^{\mu\nu} \tilde{F}_{\mu\nu}$ where N is the number of massless quarks. From this it is possible to add to the lagrangian the CP violating term: $\mathcal{L} = \theta \frac{g^2 N}{32\pi^2} F^{\mu\nu} \tilde{F}_{\mu\nu}$, which produces an electric dipole moment for the neutron
 $d_n = e(m_q/m_n)\theta \approx 10^{-16}\theta \text{e-cm}$. The current bound from PSI collaboration set $d_n < 2.9 \times 10^{26} \text{e-cm}$. Why θ is so small?, this is the strong CP problem.

Peccei-Quinn Solution

- Peccei and Quinn suggested that the SM has an additional $U(1)_{PC}$ chiral (global) symmetry which drives $\theta \rightarrow 0$. This global $U(1)_{PQ}$ symmetry was named after Roberto Peccei and Helen Quinn.

What is our work about?

Our aim is to use the $U(1)_{PQ}$ symmetry to generate quark textures motivated from the data. In particular, we are interested in the hermitian textures

$$M^U = \begin{pmatrix} 0 & 0 & |C_u|e^{i\phi_{C_u}} \\ 0 & A_u & |B_u|e^{i\phi_{B_u}} \\ |C_u|e^{-i\phi_{C_u}} & |B_u|e^{-i\phi_{B_u}} & D_u \end{pmatrix}, \quad (1)$$

$$M^D = \begin{pmatrix} 0 & |C_d| & 0 \\ |C_d| & 0 & |B_d| \\ 0 & |B_d| & A_d \end{pmatrix},$$

The PQ charges of the SM fermions are:

Particles	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$Q_{PQ}(i=1)$	$Q_{PQ}(i=2)$	$Q_{PQ}(i=3)$	$U(1)_{PQ}$
q_{Li}	1/2	3	2	1/6	$-2s_1 + 2s_2 + \alpha$	$-s_1 + s_2 + \alpha$	α	x_{q_i}
u_{Ri}	1/2	3	1	2/3	$s_1 + \alpha$	$s_2 + \alpha$	$-s_1 + 2s_2 + \alpha$	x_{u_i}
d_{Ri}	1/2	3	1	-1/3	$2s_1 - 3s_2 + \alpha$	$s_1 - 2s_2 + \alpha$	$-s_2 + \alpha$	x_{d_i}
ℓ_{Li}	1/2	1	2	-1/2	$-2s_1 + 2s_2 + \alpha'$	$-s_1 + s_2 + \alpha'$	α'	x_{ℓ_i}
e_{Ri}	1/2	1	1	-1	$2s_1 - 3s_2 + \alpha'$	$s_1 - 2s_2 + \alpha'$	$-s_2 + \alpha'$	x_{e_i}
ν_{Ri}	1/2	1	1	0	$-4s_1 + 5s_2 + \alpha'$	$-s_1 + 2s_2 + \alpha'$	$s_2 + \alpha'$	x_{ν_i}

Table: The columns 6-8 are the PQ (Q_{PQ}) charges for the SM quarks in each family. The subindex $i = 1, 2, 3$ stands for the family number in the interaction basis. The parameters s_1, s_2 and α are reals. The normalized charges are $\hat{s}_{1,2} = \frac{9}{N} s_{1,2}$ and additionally $\hat{s}_2 = (\epsilon + \hat{s}_1)$. The heavy quark PQ charges satisfy $x_L - x_R = N(1 - \epsilon)$, with $\epsilon \neq 0$.

In order to generate these matrices at least 4 higgs doublets are needed.

Particles	Spin	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	Q_{PQ}	$U(1)_{PQ}$
Φ_1	0	1	2	1/2	s_1	x_{ϕ_1}
Φ_2	0	1	2	1/2	s_2	x_{ϕ_2}
Φ_3	0	1	2	1/2	$-s_1 + 2s_2$	x_{ϕ_3}
Φ_4	0	1	2	1/2	$-3s_1 + 4s_2$	x_{ϕ_4}
S_1	0	1	1	0	$x_{S_1} = s_1 - s_2 \neq 0$	x_S
S_2	0	1	1	0	$x_{S_2} = x_R - x_L \neq 0$	x_S
Q_L	1/2	3	0	0	$x_{Q_L} - x_{Q_R} \neq 0$	x_{Q_L}
Q_R	1/2	3	0	0		x_{Q_R}

Table: Beyond SM scalar and fermion fields and their respective PQ charges. The parameters s_1, s_2 and α are reals, with $s_1 \neq s_2$. The parameters s_1, s_2 and α are reals. The normalized charges are $\hat{s}_{1,2} = \frac{9}{N}s_{1,2}$ and additionally $\hat{s}_2 = (\epsilon + \hat{s}_1)$. The heavy quark PQ charges satisfy $x_L - x_R = N(1 - \epsilon)$, with $\epsilon \neq 0$.

by choosing the VEV in a convenient way it is possible to reproduce the quark masses and the CKM mixing matrix and Yukawa couplings of order 1.

$$Y_{ij}^{U,D} \sim 1. \quad (2)$$

So, by setting various Yukawa (for quarks) couplings close to 1 (except y_{23}^{U2} , y_{23}^{D3} and y_{13}^{U1}) we obtain:

$$\hat{v}_1 = 1.71 \text{ GeV}, \quad \hat{v}_2 = 2.91 \text{ GeV}, \quad \hat{v}_3 = 174.085 \text{ GeV}, \quad \hat{v}_4 = 13.3 \text{ MeV}. \quad (3)$$

High energy Lagrangian

$$\begin{aligned}
 \mathcal{L}_{\text{LO}} \supset & (D_\mu \Phi^\alpha)^\dagger D^\mu \Phi^\alpha + \sum_\psi i \bar{\psi} \gamma^\mu D_\mu \psi + \sum_{i=1}^2 (D_\mu S_i)^\dagger D^\mu S_i \\
 & - \left(\bar{q}_{Li} y_{ij}^{D\alpha} \Phi^\alpha d_{Rj} + \bar{q}_{Li} y_{ij}^{U\alpha} \tilde{\Phi}^\alpha u_{Rj} \right. \\
 & \quad \left. + \bar{\ell}_{Li} y_{ij}^{E\alpha} \Phi^\alpha e_{Rj} + \bar{\ell}_{Li} y_{ij}^{N\alpha} \tilde{\Phi}^\alpha \nu_{Rj} + \text{h.c} \right) \\
 & + (\lambda_Q \bar{Q}_R Q_L S_2 + \text{h.c}) - V(\Phi, S_1, S_2), \tag{4}
 \end{aligned}$$

Scalar Potential

$$\begin{aligned}
 V(\Phi, S_i) = & \sum_{i=1}^4 \mu_i^2 \Phi_i^\dagger \Phi_i + \sum_{k=1}^2 \mu_{S_k}^2 S_k^* S_k + \sum_{i=1}^4 \lambda_i (\Phi_i^\dagger \Phi_i)^2 \\
 & + \sum_{k=1}^2 \lambda_{S_k} (S_k^* S_k)^2 + \sum_{i=1}^4 \sum_{k=1}^2 \lambda_{iS_k} (\Phi_i^\dagger \Phi_i) (S_k^* S_k) \\
 & + \underbrace{\sum_{i,j=1}^4}_{i < j} \left(\lambda_{ij} (\Phi_i^\dagger \Phi_i) (\Phi_j^\dagger \Phi_j) + J_{ij} (\Phi_i^\dagger \Phi_j) (\Phi_j^\dagger \Phi_i) \right) \\
 & + \lambda_{S_1 S_2} (S_1^* S_1) (S_2^* S_2) \\
 & + K_1 ((\Phi_1^\dagger \Phi_2) (\Phi_3^\dagger \Phi_2) + h.c.) \\
 & + K_2 ((\Phi_3^\dagger \Phi_4) (\Phi_3^\dagger \Phi_1) + h.c.) \\
 & + F_1 ((\Phi_2^\dagger \Phi_3) S_1 + h.c.) \\
 & + F_2 ((\Phi_1^\dagger \Phi_2) S_1 + h.c.) \\
 & + \frac{1}{2} \left(m_{\zeta S_2} \right)_{\text{SB}}^2 \zeta_{S_2}^2 + \frac{1}{2} \left(m_{\xi S_2} \right)_{\text{SB}}^2 \xi_{S_2}^2. \tag{5}
 \end{aligned}$$

It is possible to obtain dimension five effective lagrangians by means of the non-linear transformation

$$\begin{aligned} S^i &\longrightarrow e^{i \frac{x_{S^i}}{\Lambda} a} S^i, \\ \Phi^\alpha &\longrightarrow e^{i \frac{x_{\Phi^\alpha}}{\Lambda} a} \Phi^\alpha, \\ \psi_L &\longrightarrow e^{i \frac{x_{\psi_L}}{\Lambda} a} \psi_L, \\ \psi_R &\longrightarrow e^{i \frac{x_{\psi_R}}{\Lambda} a} \psi_R, \end{aligned} \tag{6}$$

Effective quark-axion interaction vertex.

from \mathcal{L}_{K^ψ} we obtain the flavour-violating derivative couplings:

$$\Delta \mathcal{L}_{K^D} = -\partial_\mu a \bar{d}_i \gamma^\mu \left(g_{af_i f_j}^V + \gamma^5 g_{af_i f_j}^A \right) d_j, \quad (7)$$

where;

$$g_{ad_i d_j}^{V,A} = \frac{1}{2f_a c_3^{\text{eff}}} \Delta_{V,A}^{Dij}, \quad (8)$$

In this expression we made the substitution $\Lambda = f_a c_3^{\text{eff}}$. The axial and vector couplings are:

$$\Delta_{V,A}^{Dij} = \Delta_{RR}^{Dij}(d) \pm \Delta_{LL}^{Dij}(q), \quad (9)$$

with $\Delta_{LL}^{Fij}(q) = \left(U_L^D \ x_q \ U_L^{D\dagger} \right)^{ij}$ and $\Delta_{RR}^{Fij}(d) = \left(U_R^D \ x_d \ U_R^{D\dagger} \right)^{ij}$

Constraints from Semileptonic decays

it is shown that the decay widths of pseudoscalar $K^\pm(B)$ mesons into an axion and a charged pion (vector K^*) are given by

$$\begin{aligned}\Gamma(K^\pm \rightarrow \pi^\pm a) &= \frac{m_K^3}{16\pi} \left(1 - \frac{m_\pi^2}{m_K^2}\right)^2 \lambda_{K\pi a}^{1/2} f_0^2(m_a^2) |g_{ads}^V|^2, \\ \Gamma(B \rightarrow K^* a) &= \frac{m_B^3}{16\pi} \lambda_{BK^* a}^{3/2} A_0^2(m_a^2) |g_{asb}^A|^2,\end{aligned}\quad (10)$$

where $\lambda_{Mma} = \left(1 - \frac{(m_a+m)^2}{M^2}\right) \left(1 - \frac{(m_a-m)^2}{M^2}\right)$

Constraints from Semileptonic decays

Collaboration	Upper bound	
E949+E787 [1, 2]	$\mathcal{B}(K^+ \rightarrow \pi^+ a) < 0.73 \times 10^{-10}$	
CLEO [3]	$\mathcal{B}(B^\pm \rightarrow \pi^\pm a) < 4.9 \times 10^{-5}$	(11)
CLEO [3]	$\mathcal{B}(B^\pm \rightarrow K^\pm a) < 4.9 \times 10^{-5}$	
BELLE [4]	$\mathcal{B}(B^\pm \rightarrow \rho^\pm a) < 21.3 \times 10^{-5}$	
BELLE [4]	$\mathcal{B}(B^\pm \rightarrow K^{*\pm} a) < 4.0 \times 10^{-5}$	

Table: These inequalities come from the window for new physics in the branching ratio uncertainty of the meson decay in a pair $\bar{\nu}\nu$.

Frame Title




Figure: Tree level diagram contribution to the FCNC processes $K^\pm \rightarrow \pi^\pm a$ and $B^\pm \rightarrow K^{*\pm} a$.

Frame Title




Figure: Allowed regions for semileptonic meson decays. We use the relation $m_a \approx 0.5m_\pi \frac{f_\pi}{f_a} \sim 0.5 \frac{m_\pi^2}{f_a}$ between the axion mass and the decay constant f_a .

Frame Title




Figure: The excluded parameter space by various experiments corresponds to the colored regions, the dashed-lines correspond to the projected bounds of coming experiments looking for axion signals, the blue region corresponds to the parameter space scanned by our model in the interval $-1 \leq \epsilon \leq 1$.

S_1 and S_2 potential.

$$\begin{aligned}
 V(\Phi, S_i) = & \sum_{k=1}^2 \mu_{s_k}^2 S_k^* S_k \\
 + & \sum_{k=1}^2 \lambda_{s_k} (S_k^* S_k)^2 + \sum_{i=1}^4 \sum_{k=1}^2 \lambda_{is_k} (\Phi_i^\dagger \Phi_i) (S_k^* S_k) \\
 + & \lambda_{s_1 s_2} (S_1^* S_1) (S_2^* S_2) \\
 + & F_1 \left((\Phi_2^\dagger \Phi_3) S_1 + h.c. \right) \\
 + & F_2 \left((\Phi_1^\dagger \Phi_2) S_1 + h.c. \right) \\
 + & \frac{1}{2} \left(m_{\zeta_{S_2}} \right)_{SB}^2 \zeta_{S_2}^2 + \frac{1}{2} \left(m_{\xi_{S_2}} \right)_{SB}^2 \xi_{S_2}^2. \tag{12}
 \end{aligned}$$

In these expressions $S_i = \frac{v_{S_i} + \xi_{S_i} + i\zeta_{S_i}}{\sqrt{2}}$; $i = 1, 2$.

dark matter connexion




Figure: The scalar potential $V(\phi_\alpha, S_1, S_2)$ is invariant under the symmetry $S_2 \rightarrow S_2^\dagger$ (which is equivalent to a Z_2 symmetry), but this symmetry is broken by the interaction term $\lambda_Q \bar{Q}_R Q_L S_2 + \text{h.c.}$. In fact, from this interaction, it is also possible to generate, at one loop, a mass term for the CP-odd field $\frac{1}{2} (m_{\zeta_{S_2}})_{\text{SB}}^2 \zeta_{S_2}^2$ in the effective Weinberg-Coleman potential.

Conclusions

- In this work we have proposed a PQ symmetry that gives rise to quark mass matrices with five texture-zeros. This texture can adjust in a non-trivial way the six masses of the quarks and the three CKM mixing angles and the CP violating phase.
- Since in our model the PQ charges are non-universal there are FCNC at the tree level. We calculated the tree level FCNC couplings from the effective interaction Lagrangian between the kinetic term of the quarks and the axion, these couplings are well known in the literature.
- The model has as a candidate for dark matter a pseudo-scalar field, which is identical to that of the Complex Singlet Extension of the Standard Model.

Scalar Potential

1. “Appears and disappears.” Burchell et al (1998)
2. Appears and disappears.

Scalar Potential

3. Reappears

- Appears;
- Appears
- Appears
- Appears

Descriptive statistics

Add an image or table.

Name	Turn	Height
Juan	1	1.9
Jose	2	1.7
Michael	3	1.95

Model

$$Y_i = \beta_0 + \beta_1 X_i + \beta_2 X_{2i} + \beta_3 Other_i + \gamma S_i + u_i \quad (13)$$

Y: Externality

- Var1.
- Var2.
- Var3.
- Var4.

Var7

Var8: Socioeconomic characteristics

- Var8.1
- Var8.2.
- Var8.3.
- Median income.
- Median home value.

S: State dummies


Outline

- 1 Strong CP problem
- 2 The PQ solution.
- 3 The hierarchy problem
- 4 The model particle content.
- 5 Low energy constraints
- 6 A dark matter candidate
- 7 Conclusions

Frame title

Name	Turn	Height
Juan	1	1.9
Jose	2	1.7
Michael	3	1.95

ReSULT



Result

Name	Turn	Height
Juan	1	1.9
Jose	2	1.7
Michael	3	1.95

Discussion

1. Comment1
2. Comment2.
3. Comment3.
4. Comment4.

Frame Title

-  S. Adler *et al.*, "Measurement of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ branching ratio," *Phys. Rev. D*, vol. 77, p. 052003, 2008.
-  A. Artamonov *et al.*, "New measurement of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ branching ratio," *Phys. Rev. Lett.*, vol. 101, p. 191802, 2008.
-  R. Ammar *et al.*, "Search for the famililon via $B^+ \rightarrow \pi^+ X_0$, $B^+ \rightarrow K^+ X_0$, and $B^0 \rightarrow K^0(S) X_0$ decays," *Phys. Rev. Lett.*, vol. 87, p. 271801, 2001.
-  O. Lutz *et al.*, "Search for $B \rightarrow h^{(*)} \nu \bar{\nu}$ with the full Belle $\Upsilon(4S)$ data sample," *Phys. Rev. D*, vol. 87, no. 11, p. 111103, 2013.

Basabendu Barman

Centro de Investigaciones

Universidad Antonio Nariño

Bogotá, Colombia

basabendu88barman@gmail.com

Bogotá, June 01,2022

To whom it may concern,

We hereby certify that Dr. Eduardo Rojas has attended the **Workshop Materia Oscura en Colombia** (MOCa 2022) (<https://indico.cern.ch/event/1113969/>), held between May 31 to June 01, 2022. Additionally, we certify that he presented the talk “*A dark matter connection in a flavored axion model*”.

Kind regards,

Basabendu Barman

Basabendu Barman, PhD