

San Juan de Pasto 7 de diciembre de 2021

Señores:

ASAMBLEA DE PROFESORES

Departamento de física

Universidad de Nariño

Cordial saludo

Por medio de la presente informo de las actividades desarrolladas durante mi comisión académica en el evento **6th Colombian Meeting on High Energy Physics (COMHEP)** que se realizó en la ciudad de Santa Marta del 29 de noviembre al 3 de diciembre del presente año.

- Presenté la charla: Faddeev-Jackiw quantization of Proca's Electrodynamics on the null-plane.
- Participé de las actividades realizadas en esta conferencia, que incluyeron: charlas principales y sesiones paralelas.

Agradezco de antemano su atención

Atentamente



German Enrique Ramos Zambrano.

Docente Tiempo Completo

Departamento de Física

Faddeev-Jackiw quantization of Proca's Electrodynamics on the null-plane

German Enrique Ramos Zambrano - Eduardo Rojas
B. M. Pimentel

Departamento de Física - Universidad de Nariño
IFT - UNESP

The effective gauge invariance Lagrangian density which describe the Proca field is defined by:

$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \frac{1}{2}M^2 \left[A_\mu + \frac{1}{e}\partial_\mu\theta \right]^2, \quad (1)$$

where $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$. The Lagrangian above is invariant by the following transformations,

$$A_\mu(x) \rightarrow A_\mu(x) + \partial_\mu\Lambda(x) \quad , \quad \theta(x) \rightarrow \theta(x) - e\Lambda(x), \quad (2)$$

From (1) it is easy to write the first-order Lagrangian by introducing the momentum π^μ and p_θ with respect to the fields A_μ and θ , respectively,

$$\pi^\mu = \frac{\partial \mathcal{L}}{\partial (\partial_+ A_\mu)} = F^{\mu+} \quad , \quad p_\theta = \frac{\partial \mathcal{L}}{\partial (\partial_+ \theta)} = \frac{M^2}{e} \left[A_- + \frac{1}{e} \partial_- \theta \right]. \quad (3)$$

from where, the starting Lagrangian density is written in first order as follow:

$$\mathcal{L}^{(0)} = \pi^- \partial_+ A_- + \pi^a \partial_+ A_a + p_\theta \partial_+ \theta - \mathcal{H}^{(0)} \quad (4)$$

where the zero iterated symplectic potential has the following form:

$$\begin{aligned}\mathcal{H}^{(0)} = & \frac{1}{2}\pi^-\pi^- + \pi^-\partial_-A_+ + \pi^a\partial_aA_+ + \frac{1}{4}F_{ab}F_{ab} - eA_+p_\theta \\ & - \frac{1}{2}M^2\left(A_a + \frac{1}{e}\partial_a\theta\right)\left(A_a + \frac{1}{e}\partial_a\theta\right).\end{aligned}\quad (5)$$

The initial set of symplectic variables is given by the set

$\xi_k^{(0)} = (A_-, A_a, \theta, \pi^-, \pi^a, p_\theta, A_+)$. We have from (4) the canonical momenta,

$$\begin{aligned}K_{A_-}^{(0)} &= \pi^-, \quad K_{A_a}^{(0)} = \pi^a, \quad K_\theta^{(0)} = p_\theta \\ K_{\pi^-}^{(0)} &= 0, \quad K_{\pi^a}^{(0)} = 0, \quad K_{p_\theta}^{(0)} = 0, \quad K_{A_+}^{(0)} = 0.\end{aligned}\quad (6)$$

The zero iterated symplectic two-form matrix is defined by,

$$M_{AB}^{(0)}(\mathbf{x}, \mathbf{y}) = \frac{\delta K_B^{(0)}(\mathbf{y})}{\delta \xi_A^{(0)}(\mathbf{x})} - \frac{\delta K_A^{(0)}(\mathbf{x})}{\delta \xi_B^{(0)}(\mathbf{y})}. \quad (7)$$

with the components

$$M_{AB}^{(0)}(\mathbf{x}, \mathbf{y}) = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\delta_b^a & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \delta_b^a & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \delta^3(\mathbf{x} - \mathbf{y}). \quad (8)$$

The symplectic matrix is singular and it has a zero mode

$$\tilde{v}^{1(0)}(\mathbf{x}) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & v(\mathbf{x}) \end{pmatrix}, \quad (9)$$

where $v(\mathbf{x})$ is an arbitrary function. From this nontrivial zero-mode, we have the following constraint,

$$\begin{aligned} \Omega_1^{(0)} &= \int d^3x v(\mathbf{x}) \frac{\delta}{\delta \xi_{A+}(\mathbf{x})} \int d^3y \mathcal{H}^{(0)}(y) \\ &= - \int d^3x v(\mathbf{x}) \left[\partial_-^x \pi^-(x) + \partial_a^x \pi^a(x) + e p_\theta(x) \right] (10) \\ &= 0. \end{aligned}$$

The constraint is evaluated from (10) to be,

$$\Omega_1^{(0)} = \partial_-^x \pi^-(x) + \partial_a^x \pi^a(x) + e p_\theta(x) = 0. \quad (11)$$

The constraint (11) is introduced in the Lagrangian density, thus, the first iterated Lagrangian density is written as,

$$\mathcal{L}^{(1)} = \pi^- \partial_+ A_- + \pi^a \partial_+ A_a + p_\theta \partial_+ \theta + \Omega_3^{(0)} \dot{\beta} - \mathcal{H}^{(1)} \quad (12)$$

where the first iterated symplectic potential is

$$\begin{aligned} \mathcal{H}^{(1)} &= \mathcal{H}_{\Omega_1^{(0)}=0}^{(0)} = \frac{1}{2} \pi^- \pi^- + \frac{1}{4} F_{ab} F_{ab} \\ &\quad - \frac{1}{2} M^2 \left(A_a + \frac{1}{e} \partial_a \theta \right) \left(A_a + \frac{1}{e} \partial_a \theta \right). \end{aligned} \quad (13)$$

We enlarged the space defined by $\xi_k^{(1)} = (A_-, A_a, \theta, \pi^-, \pi^a, p_\theta, \beta)$.

The first iterated symplectic matrix is written as,

$$M_{AB}^{(1)}(\mathbf{x}, \mathbf{y}) = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\delta_b^a & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & -\partial_-^x \\ 0 & \delta_b^a & 0 & 0 & 0 & 0 & -\partial_a^x \\ 0 & 0 & 1 & 0 & 0 & 0 & e \\ 0 & 0 & 0 & -\partial_-^x & -\partial_b^x & -e & 0 \end{pmatrix} \delta^3(\mathbf{x} - \mathbf{y}) \quad (14)$$

The modified symplectic matrix is again singular. The new constraint is identically zero.

In order to obtain a regular symplectic matrix a gauge fixing term must be added. We choose the null-plane gauge $\Theta = A_-(x) = 0$. We obtain the second iterative Lagrangian, i.e.:

$$\mathcal{L}^{(2)} = \pi^- \partial_+ A_- + \pi^a \partial_+ A_a + p_\theta \partial_+ \theta + \Omega_3^{(0)} \dot{\beta} + \Theta \dot{\eta} - \mathcal{H}^{(2)} \quad (15)$$

where

$$\begin{aligned} \mathcal{H}^{(2)} &= \mathcal{H}_{\Omega_{i1}^{(0)}, \Theta=0}^{(1)} = \frac{1}{2} \pi^- \pi^- + \frac{1}{4} F_{ab} F_{ab} \\ &\quad - \frac{1}{2} M^2 \left(A_a + \frac{1}{e} \partial_a \theta \right) \left(A_a + \frac{1}{e} \partial_a \theta \right). \end{aligned} \quad (16)$$

The new set of symplectic variable is:

$\xi_k^{(2)} = (A_-, A_a, \theta, \pi^-, \pi^a, p_\theta, \beta, \eta)$ and from (15) we determine the second-iterated symplectic two-form matrix,

$$M_{AB}^{(2)}(\mathbf{x}, \mathbf{y}) = \frac{\delta K_B^{(2)}(\mathbf{y})}{\delta \xi_A^{(2)}(\mathbf{x})} - \frac{\delta K_A^{(2)}(\mathbf{x})}{\delta \xi_B^{(2)}(\mathbf{y})}. \quad (17)$$

Since this matrix is not singular, we finally have the inverse matrix after a laborious calculation. From this relations and

$$\left\{ \xi^{(2)A}(\mathbf{x}), \xi^{(2)B}(\mathbf{y}) \right\} = \left[M_{AB}^{(2)}(\mathbf{x}, \mathbf{y}) \right]^{-1}, \quad (18)$$

we immediately identify the generalized brackets as follow:

$$\left\{ A_a(x), \pi^-(y) \right\} = -\frac{\partial_a^x}{\partial_-^x} \delta^3(\mathbf{x} - \mathbf{y}),$$

$$\left\{ A_a(x), \pi^b(y) \right\} = \delta_b^a \delta^3(\mathbf{x} - \mathbf{y}),$$

$$\left\{ \theta(x), \pi^-(y) \right\} = \frac{e}{\partial_-^x} \delta^3(\mathbf{x} - \mathbf{y}),$$

$$\left\{ \theta(x), p_\theta(y) \right\} = \delta^3(\mathbf{x} - \mathbf{y}).$$

- The results give us the Dirac brackets of the theory.
- The structure of these constraints is very simple.
- The potential symplectic obtained at the final stage of iterations is exactly the Hamiltonian.